

# Invited Speakers

## S6.4

### Early Nutritional Strategies

Y NOY<sup>1</sup> and Z UNI<sup>2</sup>

<sup>1</sup> Miloubar Feed Mill, MP Ashrat, Israel, <sup>2</sup>Faculty of Agriculture, Hebrew University, Israel

---

The embryonic and immediate posthatch developmental period represents a significant phase in attaining quality broiler performance at marketing. An efficient transition period from late term embryo to a viable independent chick is necessary for achieving results. Immediately post hatch birds must undergo a shift from egg and embryonic nutrients to exogenous feed. Under practical conditions many birds have access to feed only 36-48 h after hatching and during this time body weight decreases, intestine and muscle development is retarded. In order to overcome these limitations a continuous feeding process can be established which would supply nutrients to the developing embryo, feed and water to the newly hatched chick within the hatchery and a highly digestible prestarter diet at placement. In ovo feeding stimulates intestinal development by enhancing villi, increasing intestinal capacity to digest and absorb nutrients and provide a basis for muscle growth. Immediate access to feed (1 hr after clearing the shell) initiates uptake and growth processes some 24 h post ingestion compared to poultry with delayed feed intake. The enhanced growth caused by early feeding improves nutritional maturity of the bird, stimulates yolk utilization, increases intestinal development, and has long term metabolic effects. Providing highly digestible ingredients in the prestarter diet increases body weight performance at 7 d and through marketing. Together, these processes provide appropriate nutrition pre and post hatch which can accelerate GIT development, muscle growth and result in increased performance. This paper will summarize studies dealing with the different approaches to early nutritional strategies in our modern fast growing broiler.

---

Keywords: embryo, chick, in ovo feeding, early feeding

### Description of Problem

Birds are precocial and will forage for feed immediately and begin to grow, whereas holding them without feed results in body weight loss for some 24 h after poultry are fed (Moran, 1990, Pinchasov and Noy, 1993, Henderson et al. 2008). Under commercial practice eggs within a single tray will hatch over a 24-36 hr window during which time the birds which have piped are without feed. The early hatching chicks are therefore at a disadvantage because of the prolonged fasting period and potential dehydration (Tweed, 2005). Logistics within the hatchery as well as treatments such as sex determination, sorting, vaccinations, beak trimming, comb dubbing and transport to the farm involve a further holding period. During this time chicks decrease in weight at an approximate rate of 4 g per 24 h due in part to moisture loss as well as yolk and pectoral muscle utilization (Noy and Sklan, 1998 Halevy et al. 2003, Tona et al. 2003 and Careghi et al. 2005). Thus, birds are often held for 48 h or more before initial access to feed and water which can have lasting negative effects (Tarvid, 1992; Knight and Dibner, 1998; Noy and Sklan 2001; Batal and Parsons, 2002; Juul-Madsen et al. 2004). Delaying access to feed and water makes hatchlings more susceptible to pathogens (Dibner et al., 1998), increased weight loss (Noy and Sklan, 1999) and the development of critical tissues is restricted (Halevy et al. 2000; Moore et al. 2005) All this leads to irreversible damage to muscle development and retarded growth through marketing. These effects are more pronounced in the modern strains of meat-type poultry which are characterized by elevated metabolic rate compared to older strains (Tona et al. 2004).

### Relevance of early feeding strategies for today's poultry

At present broilers reach slaughter weight physiologically younger, the embryonic developmental period as well as the first week after hatching represents a larger proportion (45%) of the whole life span (Bigot et al., 2003). Thus, prehatch as well as the transitional posthatch period must be achieved efficiently. It is known that body weight is increased three to fourfold during the first week and considerable changes in gut, muscle weight and morphology are observed (Jin et al., 1998). Meat type broilers are capable of achieving 70g/d till 40 d. This achievement requires emphasis on early phase nutrition- "prestarter diets". Using prestarter diets assumes that starter diets are somewhat inefficient in providing balanced nutrients.

# Invited Speakers

Consequently, modern broiler strains require reevaluating nutrient provisions. Each gram of additional weight at 7 d of age translates to 5g of extra body weight at 49d (Leeson and Summers, 2005). Therefore, delayed feeding in the first few days of life reduces final BW (Noy and Sklan, 1999, Kidd et al, 2007) and it probably affects immunological capacities (Dibner et al., 1998). Early post hatching feeding is recommended to reduce these effects.

In order to develop efficient early feeding techniques it is necessary to understand the physiological and metabolic changes which occur in the pre-post hatch period.

## **Physiological Changes occurring during the pre- post hatch period**

The pre- hatch period (last phase of incubation) is characterized by oral consumption of the amnion by the embryo, accumulation of glycogen reserves in muscle and liver tissues and glycogenolysis, initiation of pulmonary respiration, abdominal internalization of remaining yolk, shell pipping, and emergence (Christensen and Biellier, 1982; Donaldson and Christensen, 1991; Donaldson *et al.*, 1991; Christensen *et al.*, 2001; Moran, 2007). During this time frame dramatic physiological and metabolic changes occur, and any disturbances at this stage may markedly affect embryonic survival and subsequent performance (Christensen *et al.*, 1999; Collin *et al.*, 2007; Leksrisonpong *et al.*, 2007). . Some of the important metabolic pathways prior to hatch are described in a recent review (de Oleveira et al 2008) which emphasizes the liver, pectoral muscle, hatching muscle and intestine, as most affected by changes toward hatch.

One of the major physiological processes during pre-natal development is the maintenance of glucose homeostasis. The glycogen reserves are withdrawn as embryos go through the hatching process. Researchers suggested insufficient glycogen and albumen forces the embryo to mobilize more muscle protein for gluconeogenesis thereby reducing early growth and development (Vieira and Moran 1999ab). The glycogen reserves begin to be replenished when the newly hatched chick has full access to feed (Moran 2007). Another significant physiological process occurs in the intestine. Exploration of this process shows that towards the end of incubation GIT undergoes extensive morphological, cellular and molecular changes. Research in broiler embryos has shown that during the last days of incubation intestinal weight relative to embryonic weight increases from 1.4% at 17E to 3.4% at hatch (Uni et al 2003). Activity and RNA expression of brush-border enzymes which digest disaccharides (sucrase-isomaltase), small peptides (aminopeptidase) and major transporters (sodium-glucose transporter and ATPase) begin to increase a few days before hatch and this process continues on day of hatch and thereafter [Gilbert, 2007; Uni et al 2003]. In the newly hatched chick the small intestinal mucosa appears to be immature. This is reflected in the organization and establishment of the crypt region, several-fold increase in villus height and area and the maturation of enterocytes and goblet cells (Uni 2006).

The immediate post-hatch period is critical for intestinal morphological development in order to digest feed and assimilate nutrients. Intestinal growth occurs in delayed fed chicks however to a significantly lesser extent than in early fed birds (Noy and Sklan, 1999 Bigot et al. 2003). This indicates a preferential intestinal growth immediately post hatch. Decreased intestinal development in chicks fasted for 36 to 48 h post-hatch was reflected in the decreased enterocyte number, crypt size, the number of crypts per villus, crypt proliferation, villus area, rate of enterocyte migration, goblet-cell size and mucin dynamics (Uni et al 2002; Geyra et al 2001).

Therefore, the sooner the GIT achieves functional capacity, the quicker the chick can utilize dietary nutrients and replenish its depleted energy status efficiently thus achieving genetic growth potential, while resisting infectious and metabolic diseases.

## **Addressing the problem -When can "early feeding" begin?**

A broad spectrum of knowledge has shown that chicks are capable of utilizing nutrients prior to hatch, during the hatching process and in transport boxes. Our proposed concept is to set aside the "traditional feeding model" of starter, grower finisher on farm feeding and provide a "nutrient link" between the embryonic and on farm growth phases. The "nutrient link" may include nutrient solutions injected into the fertile egg, feed and or water in hatcher, and during transport as well as specially designed prestarter feeds.

We will present these nutritional possibilities in the following sections;

# Invited Speakers

## *"Feeding the embryo" (In ovo feeding)*

In order to overcome the described physiological limitations and to improve intestinal functionality and nutritional status of hatchlings, a methodology for "feeding the embryo" (in-ovo feeding) was developed. A method of inserting nutrient solutions into the embryonic amniotic fluid was created for poultry and patented (Uni and Ferket 2003). The method makes use of the knowledge that neonatal birds naturally consume the amniotic fluids towards hatch (Romanoff 1960). Therefore, addition of a nutrient solution to the embryonic amniotic fluid delivers essential nutrients into the embryo intestine.

Many potential nutrient supplements can be included in the in-ovo feeding solution. Carbohydrates can be used as a source for glucose, which is crucial for the hatching process and hatchling development (Moran, 1985).  $\text{Na}^+$  and  $\text{Cl}^-$  ions play a major role in the activity of apical and basolateral transporters and in the absorption of glucose and amino acids.  $\beta$ -hydroxy- $\beta$ -methylbutyrate (HMB), a leucine metabolite which affect muscle satellite cells and increases carcass yield (Nissen et al 1994; Kornasio et al 2009), is a good candidate for the in-ovo feeding solution, as are minerals and vitamins which support the development of skeletal, immune and digestive systems in chickens.

Studies have shown that the administration of 1 ml of in-ovo feeding solution including dextrin (as a source of carbohydrates),  $\text{Na}^+$ ,  $\text{Cl}^-$ , zinc-methionine and HMB leads to increased total liver glycogen in the pre-hatch period (by 75% on the day before hatch, and by 47% on day of hatch) and markedly enhances enteric development (Tako et al 2004; Tako et al 2005; Smirnov et al 2006). Morphological evaluation of enteric sections from embryos and hatchlings revealed a significant acceleration of development 48 h after in-ovo feeding relative to non-injected controls. The in-ovo-fed birds exhibited increased pancreatic capacity for carbohydrate digestion, increased villus dimensions, higher levels of mRNA expression and activity of brush-border digestive enzymes and transporters (leucine-aminopeptidase, sucrase-isomaltase, sodium-glucose co-transporter,  $\text{Na}^+\text{K}^+$ -ATPase, zinc transporter). It can be concluded that at the time of hatch, the small intestine of in-ovo-fed birds is at a functional stage similar to that in conventionally fed 2-day-old chicks. Several experiments with offspring from young maternal flocks demonstrated that in-ovo feeding increases hatching weights by 5% over controls and elevates relative breast-muscle size (calculated as % of BW) by 6%. These weight advantages were sustained throughout the experiments (d 25). Commercial practice can be adapted to use in-ovo feeding methodology by using existing automated systems for in-ovo vaccination. Adaptation of injection machines to inject 0.6 ml to broiler and turkey embryos at 18E and 24E respectively were done by Embrex-Pfizer and in ovo feeding solutions were formulated and tested.

In-ovo feeding is expected to yield several advantages, among them reduced post-hatch mortality and morbidity; greater efficiency of feed-nutrient utilization at an early age; improved immune response to enteric antigens; reduced incidence of developmental skeletal disorders; and increased muscle development and breast-meat yield.

## *Immediate post hatch feed (Feed and/or Water in hatcher and in transport)*

A series of studies provided chicks at hatching (within one h of clearing the shell) with nutrients and examined the effect on BW until marketing d as compared to chicks held for 48 h. One such study addressed the effect of the form of the feed presented comparing solid feed, a liquid nutrient supplement or water to birds held for 48 h on BW of chicks through marketing (Noy and Sklan, 1997). Provision of caloric nutrients in solid or liquid form produced a considerable increase in BW which was maximal between 4-8 d and then decreased. Supplying water alone also resulted in an increase in BW, but this effect was smaller than that of feed and was no longer apparent after 8 d. Recent research by B. D. Farichild et al, (2006) concurs that providing chicks with water during holding increases placement weight but does not influence broiler performance after first 2 weeks of grow-out. At marketing, all birds with early access to nutrient or nutrient solutions were 8-10% heavier than those held for 48 hrs or watered birds. The cumulative feed efficiency through marketing was not changed by early nutrition, whereas the percentage of breast meat was increased by 7-9% in all fed birds (Noy and Sklan, 1998b).

Since provision of nutrients enhanced growth, the effect of applying specific materials by gavaging birds at hatch was examined using glucose, starch, protein, fat or mixtures at hatch and then returning the birds to the incubation trays. Gavaging with all nutrients enhanced BW posthatch, although glucose produced the lowest and most transient response (Moran, 1990, Pinchasov and Noy, 1993, Noy and Sklan, 1997). In an additional trial the effect of administering a single immediate posthatch intubation of nutrients to offspring of either mature (65 wks) breeder flock or young maternal flock (28 wk) was compared to held birds. Intubation significantly improved BW performance especially in offspring from young maternal flocks (Noy and Pinchasov, 1993).

# Invited Speakers

Additional techniques to reduce the adverse effects of delayed feeding have recently been researched. These include the usage of supplements known as "early feeding supplements" whose aim is to provide neonate with additional nutrient sources prior to full access to feed and water. Various commercial methods of immediate feeding have been investigated. Using Earlybird™ showed that significant increases in body weight (2.7%) can be seen at 21 and 42 d of age (Henderson et. al. 2008). Oasis™, decreased body weight loss in the 48 h post hatch period and treated chicks maintained this advantage through 21 d and 39 d of age with a higher breast yield (Noy and Sklan, 1997. Batal and Parson, 2002). Additional research involved usage of Oasis™ on female broiler breeders and indicated increased initial body weight growth (Boersma et al. 2003). Thus providing an early feeding supplement may help decrease negative effect of delayed feeding.

## *Specially Designed "Prestarter Feeds"*

On the farm feed is provided for the newly arrived chicks. However, optimal nutrition during this period must take into account the contribution of the yolk nutrients, and the ability and necessity to effectively utilize exogenous feed by the immature developing intestine. Thus, formulating prestarter diets for quick growing meat type broilers requires use of "highly digestible" ingredients. The aim of the prestarter diet is to precondition the chick such that it can digest complex substrates or provide more digestible substrate until enzyme development matures. Chicks provided with highly digestible ingredients have achieved a 7 d BW of 200g as compared with 160-170g with conventional corn-soybean diets. This increase BW is maintained through marketing (Leeson, 2008).

**In summary:** "Early" access to feed stimulates body weight growth and maintains this advantage through marketing. Early nutritional strategies offer the promise of sustaining progress in production efficiency and welfare of commercial poultry. A better understanding of the transition period from embryo to chick will further develop this "nutrient link."

## **Conclusions: Applications**

In order to achieve broiler potential we must make several unconventional changes:

- Supplying nutrients to the developing embryo before hatch
- Supplying feed within hatchery
- Developing highly digestible prestarter diets
- Integrate all of the above to create a continuous feeding process from several days prior to hatch, during hatch and till farm placement where first feed is provided.

## **References:**

- ANTHONY, N.B., DUNNINGTON, E.A.. AND SIEGEL, P.B.** (1989) Embryo growth of normal and dwarf chickens from lines selected for high and low body weight. *Archives fur Geflugelkunde*, 53: 116-122.
- BATAL, A.B. AND PARSONS C.M.** (2002) Effect of fasting versus feeding oasis after hatching on nutrient utilization in chicks. *Poultry Science* **81**: 853-859.
- BIGOT, K., MIGNON GRASTEAU S., PICARD, M. AND TESSERAUD, S.** (2003) Effects of delayed feed intake on body, intestine and muscle development in neonate broilers. *Poultry Science* 82: 781-788.
- BOERSMA, S.T., ROBINSON, F.E., RENEMA, A. AND FASENKA, G.M.** (2003) Administering Oasis™ hatching supplement prior to chick placement increases initial growth with no effect on body weight uniformity of female broiler breeders after 3 weeks of age. *Journal of Applied Research* **12**:428-434.
- CAREGHI, C., TONA, K., ONAGBESAN, O., BUYSE, J., DECUYPERE, E., AND BRUGGEMAN, V.** (2005) The effects of the spread of hatch and interaction with delayed feed access after hatch on broiler performance until seven days of age. *Poultry Science* **84**: 1314-1320.
- CHRISTENSEN, V. L., WINELAND, M. J., FASENKO, G. M. and DONALDSON, W. E.** (2001) Egg storage effects on plasma glucose and supply and demand tissue glycogen concentrations of broiler embryos. *Poultry Science* **80**:1729-1735.
- CHRISTENSEN, V. L. and BIELLIER, H. V.** (1982) Physiology of turkey embryos during pipping and hatching. IV. Thyroid function in embryos from selected hens. *Poultry Science* **61**:2482-2488.
- CHRISTENSEN, V. L., DONALDSON, W. E. and NESTOR, K. E.** (1999) Length of the plateau and pipping stages of incubation affects the physiology and survival of turkeys. *British Poultry Science* **40**:297-303.
- COLLIN, A., BERRI, C., TESSERAUD, S., RODON, F. E., SKIBA-CASSY, S., CROCHET, S., DUCLOS, M. J., RIDEAU, N., TONA, K., BUYSE, J., BRUGGEMAN, V., DECUYPERE, E.,**

# Invited Speakers

- PICARD, M. and YAHAV, S.** (2007) Effects of thermal manipulation during early and late embryogenesis on thermotolerance and breast muscle characteristics in broiler chickens. *Poultry Science* **86**:795-800.
- DIBNER, J.J., KITCHELL, M.L., ATWELL, C.A., DOWNS, A.C., AND IVEY, F.J.** (1996) The effect of dietary ingredients and age on the microscopic structure of the gastrointestinal tract in poultry *Journal Applied Poultry Research* **5**:70-77.
- DIBNER, J. J., KNIGHT, C.D., KITCHELL, M.L., ATWELL, C.A. DOWNS, A.C. AND IVEY, F.J.** (1998) Early feeding and development of the immune system in neonatal poultry. *Journal Applied Poultry Research* **7**:425-436.
- DONALDSON, W. E., and CHRISTENSEN, V. L.** (1991) Dietary carbohydrate level and glucose metabolism in turkey poults. *Comparative Biochemistry and Physiology A* **98**:347-350.
- DONALDSON, W. E., CHRISTENSEN, V. L. and KRUEGER, K. K.** (1991) Effects of stressors on blood glucose and hepatic glycogen concentrations in turkey poults. *Comparative Biochemistry and Physiology A* **100**:945-947
- DE OLIVEIRA J.E, UNI Z. AND FERKET P.R.** (2008) Important metabolic pathways in poultry embryos prior to hatch . *World Poultry Science Journal* **64**
- FAIRCHILD, B.D., NORHCUT, J.K., MAULDING , J.M., BUHR, R.J., RICHARDSON, L.J. AND COX, N.A.** (2006) Influence of water provision to chicks before placement and effects on performance and incidence of unabsorbed yolk sacs. *Journal Applied Poultry Research* **15**: 538-543.
- GEYRA, A., UNI, Z., and SKLAN, D.** (2001) Enterocyte dynamics and mucosal development in the posthatch chick. *Poultry Science* **80**:776-782.
- GILBERT, E. R., LI, H., EMMERSON, D.A., WEBB, K. E JR., WONG, E. A.** (2007) Developmental regulation of nutrient transporter and enzyme mRNA abundance in the small intestine of broilers. *Poultry Sci.* **86**(8):1739-53.
- HALEVY, O., GEYRA, A., BARAK, M., UNI, Z. AND SKLAN, D.** (2000) Early posthatch starvation decreases satellite cell proliferation and skeletal muscle growth in chicks. *Journal of Nutrition* **130**:858-864.
- HALEVY, O., NADEL, Y., BARAK, M., ROZENBOIM, I. AND SKLAN, D.** (2003) Early posthatch feeding stimulates satellite cell proliferation and skeletal muscle growth in turkey poults. *Journal of Nutrition* **133**: 1376-1382.
- HENDERSON, S.N., VICENTE, J.L., PIXLEY, C.M., HARGIS, B.M. AND TELLEZ, G.** (2008) Effect of an early nutritional supplement on broiler performance. *International Journal of Poultry Science* **7**: 211-214.
- JIN, S.H. , CORLESS, A. AND SELL, J.L.**(1998) Digestive system development in post hatch poultry. *World's Poultry Science Journal* **54**:335-345.
- JUUL-MADSEN, H.R., SU, G. AND SORENSEN, P .**(2004) Influence of early or late start of first feeding on growth and immune phenotype of broilers. *British Poultry Science* **45**:210-222.
- KIDD, M.T., TAYLOR, J.W., PAGE, C.M., LOTT, B.D., AND CHAMBLEE, T.N.** (2007) Hatchery feeding of starter diets to broiler chicks. *Journal of Applied Poultry Research* **16**: 234-239.
- KORNASIO R, RIEDERER I, BUTLER-BROWNE G, MOULY V, UNI Z, HALEVY O.** (2009) beta-hydroxy-beta-methylbutyrate (HMB) stimulates myogenic cell proliferation, differentiation and survival via the MAPK/ERK and PI3K/Akt pathways. *Biochim Biophys Acta*.
- KNIGHT, C.D. AND DIBNER, J.J.**(1998) Nutritional programming in hatchling poultry. Why a good start is important. *Poultry Digest.* **Aug/Sept.** pp:20-26.
- LEESON, S. AND SUMMERS, J.D.** (2001) *Scott's Nutrition of the Chicken.* Publ. Univ. Books, Guelph, Ontario Canada.
- LEESON, S.** (2008) Predictions for commercial poultry nutrition. *Journal of Applied Poultry Research* **17**:315-322.
- LEKSRISOMPONG, N., ROMERO-SANCHEZ, H., PLUMSTEAD, P. W., BRANNAN, K. E. and BRAKE, J.** (2007) Broiler incubation. 1. Effect of elevated temperature during late incubation on body weight and organs of chicks. *Poultry Science* **86**:2685-2691
- MOORE, D.T., FERKET, P.R. AND MOZDZIAK, P.E.** (2005) The effect of early nutrition on satellite cell dynamics in the young turkey. *Poultry Science* **84**:748-756.
- MORAN, E.T.** 1985. Digestion and absorption in fowl and events through perinatal development. *J. Nutrition* **115**(5):665-674.
- MORAN, E.T.** (2007) Nutrition of the developing embryo and hatchling. *Poultry Science* **86**:1043-1049.
- MURAKAMI, H., AKIBA, Y. AND HORIGUCHI, M.** (1992) Growth and utilization of nutrients in newly hatched chicks with or without removal of residual yolk. *Growth Development and Aging*, **56**: 75-84.
- NISSEN, S., FULLER, JR. J.C., SELL, J. FERKET, P.R. AND RIVES, D.V.** (1994) The effect of beta-hydroxy-beta-methylbutyrate on growth, mortality, and carcass qualities of broiler chickens. *Poultry Science* **73**:137-155 .

# Invited Speakers

- NOY, Y. AND SKLAN, D.** (1999a) Different types of early feeding and performance in chicks and poult. *Journal of Applied Poultry Research* **8**:16-24.
- NOY, Y. AND SKLAN, D.** (1999b) Energy utilization in newly hatched chicks. *Poultry Science* **78**:1750-1756.
- SMIRNOV, A., TAKO E., FERKET, P.R. AND UNI. Z.** (2006) Mucin gene expression and mucin content in the chicken intestinal goblet cells are affected by in ovo feeding of carbohydrates. *Poultry Science* **85**:669-673.
- TAKO, E., FERKET, P.R. AND UNI, Z.** (2004) The effects of In Ovo feeding of carbohydrates and beta-hydroxy-beta-methylbutyrate on the development of chicken intestine. *Poultry Science* **83**:2023-2028.
- TAKO, E., FERKET, P.R. AND UNI, Z.**(2005). Changes in chicken intestinal zinc exporter (ZnT1) mRNA expression and small intestine functionality following an intra amniotic zinc-methionine (ZnMet) administration. *J. Nutr. Biochem.* **16**(6):339-346.
- TARVID, I.** (1992) Effect of early postnatal long term fasting on the development of peptide hydrolysis in chicks *Comparative Biochemistry Physiology* **1010**: 161-166.
- TONA, K., ONAGBESAN, O.M., JEGO, Y., KAMERS, B., DECUYPERE, E., AND BRUGGEMAN, V.** (2004) Comparison of embryo physiological parameters during incubation, chick quality, and growth performance of three lines of broiler breeders differing in genetic composition and growth rate. *Poultry Science* **83**:507-513.
- TONA, K. F. , BARNELIS, B., DE KETTELAERE, V. BRUGGERMAN, V. MORAES, B. M. , VUYSE , J., ONAGBESAN, O. AND DECUYPERE, E.** (2003) Effects of egg storage time on spread of hatch, chick quality and chick juvenile growth. *Poultry Science* **82**: 736-741.
- TWEED, S.** (2005)The hatch window. *Cobb-Vantress Technical Focus*. Vol. 2 Siloam Springs, AR.
- UNI Z.** (2006) Early development of small intestinal function. *Avian Gut Function in Health and Disease* CAB International. p 29-42.
- UNI, Z. AND FERKET P.R.** (2003) Enhancement of development of oviparous species by in ovo feeding. US patent number 6,592,878. Issued: Jul 15, 2003.
- UNI, Z., TAKO, E., GAL-GARBER, O. and SKLAN, D.** (2003) Morphological, molecular, and functional changes in the chicken small intestine of the late-term embryo. *Poultry Science* **82**:1747-1754.
- UNI, Z., SMIRNOV A. AND SKLAN D.** (2002) Pre- and posthatch development of goblet cells in the broiler small intestine: effect of delayed access to feed. *Poultry Sci.* **82**:320-327.
- VIEIRA S.L. AND MORAN E.T.** (1999a) Effect of egg origin and chick post hatch nutrition on broiler live performance and meat yields *World poultry science journal* **56**:125-142.
- VIEIRA S.L. AND MORAN E.T.** (1999b) Effect of delayed placement and used litter on broiler yields *Journal of Applied Poultry Research* **8**: 75-81.