

The effect of photoperiod on testis weight and comb area of male broiler breeders

N. C. TYLER* and R. M. GOUS

Faculty of Science and Agriculture, Discipline of Animal and Poultry Science, University of KwaZulu-Natal, P. Bag X01, Scottsville, 3209, South Africa.

*Corresponding author: millsni@ukzn.ac.za

Abstract. From the literature, it is evident that broiler breeder pullets exhibit photorefractoriness, and while it is assumed that male broiler breeders respond in a similar manner to females, it would be beneficial to determine if this is the case. This would enable lighting programmes to be designed that will ensure maximum fertility of both males and females. Male broiler breeders were housed in light-tight rooms and given constant photoperiods of 8, 10, 12, 14, 16 or 18 h/d. One or two birds were sampled from each room weekly from 71 to 232 d, and the remaining birds at 467 d, and testis weights measured. There was no significant difference between left and right testis weights, so the average was used in the analyses. Digital images of the head were recorded prior to slaughter and of four birds in each room weekly. The area of the comb was measured using image analysis software. The relationship between comb area and average testis weight was determined by regression analysis, with a logistic curve being best at fitting the data. A positive, significant ($P < 0.001$) relationship was observed with 69% of the variance in average testis weight being accounted for by comb area. Thus, comb area could provide a reasonable, non-invasive technique to assess fertility. At 165 d, average testis weights, as well as comb areas of birds not slaughtered, showed strong evidence that males exhibit juvenile photorefractoriness in a manner similar to females. Males on shorter photoperiods matured faster than those on photoperiods of 14 h and longer. As the birds aged, the effects of photorefractoriness wore off, but at the end of the experiment adult photorefractoriness was again evident, with birds on 14 h and longer having smaller average testis weights, and therefore, no longer responding to the stimulatory photoperiod.

Keywords: Photoperiod; Testis; Reproduction; Breeder; Male

Introduction

Broiler breeder pullets have been shown to exhibit both juvenile and adult photorefractoriness, a condition in which there is no photosexual response to an otherwise stimulatory daylength (Lewis *et al.*, 2003; Gous & Cherry, 2004). This condition is evident in meat-type hybrids, while selection for early maturity and high egg production in laying-type pullets has led to the disappearance of this phenomenon in these birds. It appears that the photoperiodic response of male and female meat-type hybrids is similar (Parker & McCluskey, 1965; Renden *et al.*, 1991; Lewis *et al.*, 2004); however, it would be beneficial to determine if this is the case, in order to maximise fertility of both males and females in the design of lighting programmes.

While female fertility may be quantitatively measured by oviposition, male fertility is more difficult to assess. Wilson *et al.* (1988) report a significant ($P < 0.001$), positive ($r = 0.79$) correlation between number of spermatozoa per ejaculate and testes weight. Data provided by Kirby *et al.* (1998) in Amann (1999) show a strong, positive relationship between testis weight and daily sperm production; however, because the testes are internal, this measurement is only possible with the use of ultrasound (Hofbauer & Krautwald-Junghanns, 1999). Many assessments of semen quality have been used in an attempt to quantify male fertility, such as volume (Brown & McCartney, 1983),

concentration (Anash *et al.*, 1983; Bilgili & Renden, 1984), morphology and mobility (Froman *et al.*, 1999; Froman & Feltmann, 2000; Donoghue *et al.*, 2003). However, this requires training of males, as well as the fact that collecting semen is not necessarily representative of semen quality during natural mating. Wilson *et al.* (1988) reported that 25% of the males used in their study that exhibited spermatogenic activity, based on visual appraisal of seminiferous tubular size and maturity of cell types, failed to ejaculate semen.

Spermatogenesis relies on testosterone, which can also be measured in the blood as an indicator of fertility (Schanbacher *et al.*, 1974; Rozenboim *et al.*, 1993). However, an external, highly correlated, non-invasive indicator of testis size would be a valuable measurement. Testosterone is also responsible for the development of secondary sexual characteristics, such as comb development (Zeller, 1971). There is literature to suggest that birds with large ornaments are expressing higher reproductive fitness. It has been shown that, as well as favouring new females or females with high promiscuity, males also favour females with large ornaments, by preferentially allocating sperm to these individuals, which are thought to signal superior maternal investment (Pizzari *et al.*, 2003). In males, comb size had a significant ($P < 0.05$) effect on male-female proximity (Graves *et al.*, 1985), where females exposed to unfamiliar males stayed closer to males with larger combs. Parker & Ligon (2002) found that crowing, a secondary sexual characteristic, occurred more frequently in larger-combed males, and females preferred males with larger, brighter combs (Zuk *et al.*, 1995). This paper investigates the relationship between comb size and testis weight in male broiler breeders, and uses the relationship to examine the effects of constant photoperiods on fertility.

Materials and methods

Day old breeders from the primary breeder¹ were given 48h light and thereafter placed on one of six constant photoperiods; 8, 10, 12, 14, 16 or 18h. Each photoperiod was replicated twice, starting with 50 birds in each light-tight room. Lights were set to come on at 07:00 for all treatments.

Birds received *ad-libitum* feed for the first three weeks, and thereafter were fed a commercial breeder ration to follow the growth curve recommended by the primary breeder. Each bird was weighed on a weekly basis, and feed was allocated to a room based on the average body weight of birds in that room.

Males were trained for semen collection, starting at 19 weeks.

One or two birds from each room, depending on the stocking density and mortality rate, were slaughtered weekly from 71d to 232d, and the remaining birds at 467d. Before slaughter, a digital photograph was taken of the left and right side of the bird, with a linear scale being included in each photograph. The area of the comb was measured with image analysis technology² (McGary *et al.*, 2003).

After slaughter, both the left and right testes were removed, weighed, and their dimensions recorded.

Statistical analysis

A general ANOVA of the average testis weight (left and right testis weights were not significantly different) and comb area for the six photoperiods was performed. A logistic regression of average testis weight and comb area was fitted using GenStat 6th edition.

Results and discussion

A positive, significant ($P < 0.001$) relationship between average testis weight and comb area was found, with 69% of the variance in average testis weight being accounted for by comb area (Figure 1), suggesting that comb area provides a valid indicator of testis weight. There appear to be a few outliers as comb size and testis weight increase. Thus, it appears possible that weak individuals may try to

¹ Ross Poultry Breeders

² ImageJ

signal dominance by displaying well-developed ornaments. However, a behavioural mechanism has been reported, in which dominant acting males (those with large combs and a higher incidence of crowing) show aggression towards like individuals, which allows “punishment of cheaters” (Parker & Ligon, 2002). A negative relationship between comb size and circulating lymphocytes has also been identified (Zuk *et al.*, 1995), inferring a cost to the immune system by maintenance of secondary sexual characteristics. In females, the relationship between comb size and plasma oestradiol concentration was small, but the change in comb size was a good indicator of age at sexual maturity (Joseph *et al.*, 2003).

Other non-invasive measurements are possible. Sperm penetration of the perivitelline membrane over the region of the germinal disc can be assessed (Bakst & Howarth, 1977; Wishart, 1997; Hazary *et al.*, 2000; McGary *et al.*, 2003) and is positively correlated with egg fertility ($r = 0.89$, $P < 0.0001$) (Bramwell *et al.*, 1995). Behavioural traits could also be used as a non-invasive indicator of fertility. Wing flapping, in particular, was correlated with dominance and fertility (Jones & Mench, 1991; Leonard and Zanette, 1998). However, measuring comb size would practically be easier than observing behaviour.




Figure 1 Logistic regression of average testis weight and comb area

Both juvenile and adult photorefractoriness were evident in this experiment, with photoperiod having a significant ($P < 0.001$) effect on average testis weight over the entire slaughter period. Photoperiods of 8 and 10 h resulted in significantly higher testis weights, whilst any photoperiod over 14 h reduced this weight. Average testis weights at 165 d (Figure 2) show evidence of juvenile photorefractoriness ($n=4$ for all photoperiods except $n=3$ for 18h) where birds were not responsive to the stimulatory photoperiods. Average testis weights from images of birds not slaughtered at 165d were predicted using the relationship established in Figure 1, and the only testis weights above 10 g were predicted from birds on the 8, 10 and 12 h photoperiods.




Figure 2 Ave testis weights at 165d of birds on different constant photoperiods.

As the birds aged, juvenile photorefractoriness was dissipated, and photoperiod had less impact on average testis weight by 189 d (Figure 3).




Figure 3 Average testis weights at 189d of birds on different constant photoperiods

Evidence of photorefractoriness was again observed in the adult form at the end of the experiment with birds on longer photoperiods showing a decline in average testis weight (Figure 4). Regression of the testes, especially in seasonal breeders, occurs due to regulated changes in number and viability of Sertoli cells, spermatogonia and spermatids (Thurston & Korn, 2000). Thus, longer photoperiods were no longer stimulatory.




Figure 4 Average testis weights at 467d of birds on different constant photoperiods

These results provide strong evidence that male broiler breeders exhibit both juvenile and adult photorefractoriness. The response to photoperiod appears to be similar to that of females, where maturity was most delayed in pullets maintained on constant 13 and 14h photoperiods (Lewis *et al.*, 2004). Therefore, these results suggest that the response of males and females to constant photoperiods is similar. Further research into the response to time of photostimulation is being conducted.

References

- AMANN, R.P. (1999) Lessons for the poultry industry gleaned from experiences with other commodity species. *Poult. Sci.* **78**:419-427.
- ANASH, G.A., BUCKLAND, R.B., CHAN, C.W. and TOUCHBURN, S.P. (1983) Effects of floor and cage housing of male turkeys on semen production and fertility of fresh and stored semen. *Poult. Sci.* **62**:1899-1901.
- BAKST, M.R. and HOWARTH, B. Jr. (1977) Hydrolysis of the hen's perivitelline layer by cock sperm in vitro. *Biol. Reprod.* **17**:370-379.
- BILGILLI, S.F. and RENDEN, J.A. (1984) Fluorometric determination of avian sperm viability and concentration. *Poult. Sci.* **63**:2275-2277.
- BRAMWELL, R.K., MARKS, H.L. and HOWARTH, B. (1995) Quantitative determination of spermatozoa penetration of the perivitelline layer of the hen's ovum as assessed on oviposited eggs. *Poult. Sci.* **74**:1875-1883.
- BROWN, H.B. and McCARTNEY, M.G. (1983) Effects of dietary restriction on reproductive performance of broiler breeder males. *Poult. Sci.* **62**:1885-1888.
- DONOGHUE, A.M, KIRBY, J.D., FROMAN, D.P., LERNER, S.P., CROUCH, A.N., KING, L.M., DONOGHUE, D.J. and SONSTEGARD, T.S. (2003) Field testing the influence of sperm competition based on sperm mobility in breeder turkey toms. *Br. Poult. Sci.* **44**:498-504.
- FROMAN, P. and FELTMANN, A.J. (2000) Sperm mobility: Phenotype in roosters (*Gallus domesticus*) determined by concentration of motile sperm and straight line velocity. *Biol. Reprod.* **62**:303-309.
- FROMAN, D.P., FELTMANN, A.J., RHOADS, M.L. and KIRBY, J.D. (1999) Sperm mobility: A primary determinant of fertility in the domestic fowl (*Gallus domesticus*). *Biol. Reprod.* **61**:400-405.
- GOUS, R.M. and CHERRY, P. (2004) Effects of body weight at, and lighting regimen and growth curve to, 20 weeks on laying performance in broiler breeders. *Br. Poult. Sci.* **45**:445-452.
- GRAVES, H.B., HABLE, C.P. and JENKINS, T.H. (1985) Sexual selection in *gallus*: Effects of morphology and dominance on female spatial behavior. *Behav. Proc.* **11**:189-197.

- HAZARY, R.C., STAINES, H.J. and WISHART, G.J.** (2000) Assessing the efficiency of mating in broiler breeder flocks by enumerating the spermatozoa which penetrate the inner perivitelline layer over the germinal disc. *Br. Poult. Sci.* **41**:395-400.
- HOFBAUER, H. and KRAUTWALD-JUNGHANS, M-E.** (1999) Transcutaneous ultrasonography of the avian urogenital tract. *Veterinary Radiology & Ultrasound* **40**:58-64.
- KIRBY, J.D., WASHINGTON, J., RHOADS, M.L. and KREIDER, D.L.** (1998) Impaired testis development and spermatogenesis in adult male fowl following unrestricted prepubertal growth and subsequent growth restriction. *Poult. Sci.* **77**(Suppl.1):91 (Abst.)
- LEWIS, P.D., CIACCIARIELLO, M. and GOUS, R.M.** (2003) Photorefractoriness in broiler breeders: sexual maturity and egg production evidence. *Br. Poult. Sci.* **44**:634-642.
- LEWIS, P.D., BACKHOUSE, D. and GOUS, R.M.** (2004) Constant photoperiods and sexual maturity in broiler breeder pullets. *Br. Poult. Sci.* **45**:557-560.
- JONES, M.E.J. and MENCH, J.A.** (1991) Behavioral correlates of male mating success in a multisire flock as determined by DNA fingerprinting. *Poult. Sci.* **70**:1493-1498.
- JOSEPH, N.S., ROBINSON, F.E., RENEMA, R.A. and THORSTEINSON, K.A.** (2003) Comb growth during sexual maturation in female broiler breeders. *J. Appl. Poult. Sci. Res.* **12**:7-13.
- LEONARD, M.L. and ZANETTE, L.** (1998) Female mate choice and male behaviour in domestic fowl. *Anim. Behav.* **56**:1099-1105.
- McGARY, S., ESTEVEZ, I. and BAKST, M.R.** (2003) Potential relationships between physical traits and male broiler breeder fertility. *Poult. Sci.* **82**:328-337.
- PARKER, T.H. and LIGON, J.D.** (2002) Dominant male red junglefowl (*Gallus gallus*) test the dominance status of other males. *Behav. Ecol. Sociobiol.* **53**:20-24.
- PARKER, J.E. and McCLUSKEY, W.H.** (1965) The effect of length of daily light periods on sexual development and subsequent fertilizing capacity of male chickens. *Poult. Sci.* **44**:23-27.
- PIZZARI, T., CORNWALLIS, C.K., LOVLIE, H., JAKOBSSON, S. and BIRKHEAD, T.R.** (2003) Sophisticated sperm allocation in male fowl. *Nature* **426**:70-74.
- RENDEN, J.A., OATES, S.S. and WEST, M.S.** (1991) Performance of two male broiler breeder strains raised and maintained on various constant photoschedules. *Poult. Sci.* **70**: 1602-1609.
- ROZENBOIM, I., SNAPIR, N., ARNON, E., BEN ARYEH, R., BURKE, W.H., SHARP, P., KOCK, Y. and ROBINZON, B.** (1993) Precocious puberty in tamoxifen-treated cockerels: Hypothalamic gonadotrophin-releasing hormone, prolactin, growth hormone and testosterone. *Br. Poult. Sci.* **34**:533-542.
- SCHANBACHER, B.D., GOMES, W.R. and VANDEMARK, N.L.** (1974) Diurnal rhythm in serum testosterone levels and thymidine uptake by tested in the domestic fowl. *J. Anim. Sci.* **38**:1245-1248.
- THURSTON, R.J. and KORN, N.** (2000) Spermiogenesis in commercial poultry species: Anatomy and control. *Poult. Sci.* **79**:1650-1668.
- WILSON, J.L., KRISTA, L.M., McDANIEL, G.R. and SUTTON, C.D.** (1988) Correlation of boiler breeder male semen production and testes morphology. *Poult. Sci.* **67**:660-668.
- WISHART, G.J.** (1997) Quantitative aspects of sperm:egg interaction in chickens and turkeys. *Anim. Reprod. Sci.* **48**:81-92
- ZELLER, F.J.** (1971) The effects of testosterone on dihydrotestosterone on the comb, testis, and pituitary gland of the female fowl. *J. Reprod. Fert.* **25**:125-127.
- ZUK, M., JOHNSEN, T.S. and MACLARTY, T.** (1995) Endocrine-immune interactions, ornaments and mate choice in red jungle fowl. *Proc. R. Soc. Lond. B* **260**:205-210.