



ESTABLISHING THE PSYLLID: FIELD STUDIES FOR THE BIOLOGICAL CONTROL OF JAPANESE KNOTWEED

Locations Canada, Japan, Netherlands, United Kingdom, United States

Dates 25/03/2008 - Ongoing

Summary

Japanese knotweed is highly damaging. It spreads extremely quickly, preventing native vegetation from growing and has significant impacts on infrastructure. Current control methods rely mainly on chemicals. Research, however, has identified a tiny psyllid from Japan as a suitable and safe agent to control Japanese knotweed in the UK, Canada, the Netherlands and USA. The current aim of this project is to achieve establishment and impact of the psyllid on Japanese knotweed in these countries.

The problem

Japanese knotweed (*Fallopia japonica*) is one of the most damaging invasive weeds in the UK, Europe and North America. Growing up to a metre a month, it is a problem to the construction industry whilst also out-competing native vegetation. Since it was introduced as a prized ornamental from Japan in the

mid-19th century, the species has become problematic in its entire exotic range where it has established.

In its native Asian range, this plant species evolved with more than one specialized natural enemy – from fungal diseases to insects – which help to control its growth and spread. In Europe, USA and Canada however, none of these natural enemies are present, meaning this non-native invasive knotweed is literally out of control.

Total costs for Japanese knotweed to the British economy is around £165.6 million per year. Currently, the weed can be partially controlled using chemicals, however, given the rate at which the species spreads, conventional control methods are deemed unsustainable because they are costly and/or impractical. Classical biological control could offer a more economical, sustainable and environmentally-friendly approach.

What we are doing

CABI's scientists have carried out extensive research into the biocontrol potential of an insect and a fungal pathogen since 2000 in the UK and USA, extending to Canada in 2007 and then to the Netherlands in 2019.

Field surveys revealed a number of possible natural agents associated to Japanese knotweed and from these, the *Aphalara itadori* psyllid (a small sap-sucking insect) and fungal leaf-spot pathogen, *Mycosphaerella polygoni-cuspidati*, were seen as having the highest potential as biocontrol agents.

CABI is working with the UK Government on the approved release of the specialist psyllid as a control agent for Japanese knotweed in the UK. The psyllid was approved for release in the UK in 2010 followed by mass releases during subsequent years. The releases are taking place in field trials at a number of sites in England and Wales. Part of the release protocol includes a monitoring plan, which aims to record the impact of the psyllid on Japanese knotweed, but also to detect any potential adverse effects on native plants and invertebrates.

Observations revealed that the psyllids collected from Mt. Aso, Kumamoto Prefecture, Japan, reproduced and survived over on Japanese knotweed at several release sites in the UK. However, long-term establishment and persistent overwintering have proved challenging.

Research in Canada assessed the suitability of the agents for this geographic region. Research for British Columbia's Ministry of Forests, Lands and Natural Resource Operations, as well as for the USA in partnership with the United States Department of Agriculture (USDA), has been running in parallel. The psyllid was released in Canada in 2014 and in the USA in 2021. The spread and impact will be closely monitored.

Murakami psyllid In 2019, a survey was carried out in Japan to collect better climatically-matched psyllids. Extensive and severe leaf curling damage, attributed to psyllids, was found at Murakami in the Niigata Prefecture and will continue to be investigated. The Murakami psyllid was released in 2020 in the Netherlands after approval for release by Dutch authorities. The psyllid was also released in the UK and Canada in 2021 – its spread and impact are being closely monitored.

***Mycosphaerella polygoni-cuspidati*, leaf-spot pathogen** The fungal leaf-spot pathogen, *Mycosphaerella polygoni-cuspidati*, was not successful as a classical agent due to its potential to damage other plants. However, it is now undergoing [research](#) to assess its suitability as a mycoherbicide.

Results so far

Further information on CABI's work on Japanese knotweed can be found [here](#).

Donors

Ministry of Forests, British Columbia, Canada, Cornwall Council - previous sponsor, British Waterways (Canal & River Trust) - previous sponsor, Department for Environment, Food and Rural Affairs (DEFRA) - current sponsor, Environment Agency - previous sponsor, Network Rail - previous sponsor, South West Regional Development Agency (SWRDA) - previous sponsor, Welsh Assembly Government - current sponsor

Partners

University of Reading, Agriculture and AgriFood Canada-Lethbridge Research Centre, Kyushu University

CABI Project Manager

Daisuke Kurose



<https://www.cabi.org/what-we-do/cabi-projects/>