

RICE ALLELOPATHY AND MOMILACTONE

Hisashi Kato-Noguchi

Department of Applied Biological Science, Faculty of Agriculture,
Kagawa University, Miki, Kagawa 761-0795, Japan

ABSTRACT

Rice has been extensively studied with respect to its allelopathy as part of a strategy for sustainable weed management options. All available information indicates that rice plants possibly release unknown allelochemicals into the neighbouring environments. A large number of compounds, such as phenolic acids, fatty acids, indoles and terpenes, were identified in rice root exudates and decomposing rice residues as putative allelochemicals. Among them, momilactone B may play an important role in rice defence mechanism in the rhizosphere for the competition with invading root systems of neighbouring plants.

Keywords: Allelopathy, Allelochemical, Momilactone, Root exudates, Sustainable weed management.

INTRODUCTION

Weeds cause reductions in rice yield and quality and remain one of the biggest problems in rice production. The negative impacts of commercial herbicide use on the environment make it desirable to diversify weed management options. Allelopathy is one of the options (Rimando and Duke 2003, Macías et al. 2007, Kong 2008, Tesio and Ferrero 2010).

Allelopathy is the direct influence of an organic chemical released from one living plant on the growth and development of other plants (Inderjit and Duke 2003, Belz 2007, Macías et al. 2007). Allelochemicals are such organic chemicals involved in the allelopathy (Rice 1984, Putnam and Tang 1986, Inderjit 1996). Allelochemicals can provide a competitive advantage for host-plants through suppression of soil microorganism and inhibition of the growth of competing plant species because of their antibacterial, antifungal, and growth inhibitory activities (McCully 1999, Hawes et al. 2000, Bais et al. 2004).

Rice has also been extensively studied with respect to its allelopathy as part of a strategy for sustainable weed management options. A large number of rice varieties were found to inhibit the growth of several plant species when these rice varieties were grown together with these plants under the field or/and laboratory conditions (Dilday et al. 1994, 1998, Kim et al. 1999, Olofsdotter et al. 1999; Azmi et al. 2000, Gealy et al. 2003, Seal et al. 2004a, Kim et al. 2005). These findings suggest that rice may produce and release allelochemicals into neighbouring environment, thus encouraging the exploration of allelochemicals in rice.

Many secondary compounds, such as phenolic acids, fatty acids, indoles and terpenes, were identified in rice root exudates and decomposing rice residues as putative allelochemicals (Takeuchi et al. 2001, Rimando and Duke 2003, Khanh et al. 2007).

However, these compounds are almost ubiquitous in plants and rice allelopathy can not be explained by these compounds (Olofsdotter et al. 2002b, Seal et al. 2004b). Diterpen, known as momilactone B, which are unique to rice, have been isolated (Kato-Noguchi et al. 2002). Momilactone B inhibits the growth of typical rice weeds like *Echinochloa crus-galli* and *E. colonum* at concentrations greater than 1 μM (Kato-Noguchi et al. 2008a). Rice plants secrete momilactone B from the roots into the rhizosphere over their entire life cycle (Kato-Noguchi et al. 2003b). These observations suggest that rice plants may inhibit the growth of the neighbouring plants through the secretion of momilactone B into their rhizosphere.

Rediscovery of Momilactone

About 5,000 rice seedlings, cv. Koshihikari, were hydroponically grown for 14 days in order to screen for any allelochemicals in rice root exudates. Keeping track of the biological activity, the culture solution was purified by several chromatographic fractionations and finally 2.1 mg of putative compound causing the inhibitory effect of the rice seedlings was isolated (Kato-Noguchi et al. 2002, Kato-Noguchi and Ino 2003a). The chemical structure of the inhibitor was determined from high-resolution MS, and ^1H - and ^{13}C -NMR spectral data as momilactone B. Momilactone B was later found in root exudates of other allelopathic rice cultivars, PI312777, with 5,7,4'-trihydroxy-3',5'-dimethoxyflavone and 3-isopropyl-5-acetoxycyclohexene-2-one-1 (Kong et al. 2004). In addition, another potential allelochemical momilactone A was found in rice root exudates of cv. Koshihikari (Kato-Noguchi et al. 2008b).

Momilactone A and B were first isolated from rice husks as growth inhibitors (Kato et al. 1973, Takahashi et al. 1976). Momilactone A and B were later found in rice leaves and straw as phytoalexins (Cartwright et al. 1977, 1981, Kodama et al. 1988, Lee et al. 1999). Thereafter, the function of momilactone A as a phytoalexin has been extensively studied and several lines of evidence indicate that momilactone A has an important role in rice defense system against pathogen attacks (Nojiri et al. 1996, Araki and Kurahashi 1999, Takahashi et al. 1999, Tamogami and Kodama 2000, Agrawal et al. 2002). Although the growth inhibitory activity of momilactone B was much greater than that of momilactone A (Takahashi et al. 1976, Kato et al. 1977), the efforts to find the function of momilactone B were limited.

Biological Activity of Momilactone

Momilactone A and B, respectively, inhibited the growth of *Amaranthus lividus*, *Digitaria sanguinalis* and *Poa annua* at concentrations greater than 20 ppm (ca. 60 μM) and 4 ppm (ca. 12 μM) (Chung et al. 2005). Momilactone A and B were also reported to inhibit the growth of *Echinochloa crus-galli* and *E. colonum*, which are the most noxious weeds in rice fields, at concentrations greater than 10 and 1 μM , respectively. Thus, effectiveness of momilactone B on growth inhibition is much greater than that of momilactone A. The growth inhibitory activities of momilactone B are also greater than those of momilactone A under other bioassay systems, (Takahashi et al. 1976, Kato et al. 1977, Fukuta et al. 2007, Toyomasu et al. 2008).

Momilactone A and B, respectively, inhibited root and shoot growth of rice seedlings at concentrations greater than 100 and 300 μM . IC_{50} values of momilactone A and B on rice root and shoot were not obtained because of their weak inhibitory activities against rice. The inhibitory activities of momilactone A and B, respectively, on the root and shoot growth

of rice seedlings were 1 - 2 % and 0.6 - 2 % of those on the root and shoot growth of *E. crus-galli* and *E. colonum*. Thus, the effectiveness of momilactone A and B on the growth of rice seedlings was much less than that on the growth of *E. crus-galli* and *E. colonum*. These results suggest that the toxicities of momilactone A and B to rice seedlings are much less than those to the two weed species (Kato-Noguchi et al. 2008a).

Momilactone in Rice Life Cycle

Momilactone A and B were secreted from rice plants into the rhizosphere throughout all life cycle stage of rice (Kato-Noguchi et al. 2003b, 2008a). The secretion level of momilactone A and B increased until flowering initiation, and decreased thereafter. The level of momilactone A and B at day 80 (around flowering) was 1.1 and 2.3 µg/plant/day, which was 55- and 58-fold greater than that at day 30. Although concentration of momilactone A in rice was greater than that of momilactone B, secretion level of momilactone B was greater than that of momilactone A, which suggests that momilactone B may be selectively secreted into the rhizosphere than momilactone A.

Considering the growth inhibitory activity and concentrations found in the bioassay medium, momilactone A may cause only 0.8 - 2.2% of the observed growth inhibition of *E. crus-galli* roots and shoots by rice. However, momilactone B in the medium was estimated to cause 59 - 82% of the observed growth inhibition of *E. crus-galli* roots and shoots by the rice seedlings. In addition, the concentrations of momilactone B in the medium reflected the observed differences in the growth inhibition of *E. crus-galli* by the eight rice cultivars investigated (Kato-Noguchi et al. 2010). This suggests that the allelopathic activity of rice may be primarily depend on the secretion level of momilactone B. Therefore, momilactone B may play a very important role in rice allelopathy.

Rice Allelopathy and Allelochemicals

Since the first observation of allelopathy in rice by Dilday et al. (1989), more than 16,000 rice accessions from 99 countries in the USDA-ARS germplasm collection have been screened. Of these, 412 accessions inhibited the growth of *Heteranthera limosa* and 145 accessions inhibited the growth of *Ammannia coccinea* (Dilday et al. 1994, 1998). Similar attempts have been conducted in some other countries, and a large number of rice varieties were found to inhibit the growth of several plant species when these rice varieties were grown together with these plants under field and/or laboratory conditions (Kim et al. 1999, Olofsdotter et al. 1999, Azmi et al. 2000, Gealy et al. 2003, Seal et al. 2004a, Kim et al. 2005). These findings suggest that rice may produce and secrete allelochemicals into its neighboring environments.

Although mechanisms of the exudation are not well understood, it is suggested that plants are able to secrete a wide variety of compounds from root cells by plasmalemma-derived exudation, endoplasmic-derived exudation, and proton-pumping mechanisms (Hawes et al. 2000, Bais et al. 2004). Through the root exudation of compounds, plants are able to regulate the soil microbial community in their immediate vicinity, change the chemical and physical properties of the soil, and inhibit the growth of competing plant species (MuCully 1999, Hawes et al. 2000, Bais et al. 2004). Momilactone B was secreted from rice plants into the rhizosphere throughout all life cycle stage of rice (Kato-Noguchi et al. 2003b, 2008a). Considering the inhibitory activity and the secretion level, momilactone B may play a very important role in rice defense mechanism in the rhizosphere as an allelochemical.

The use of allelopathic rice cultivars and allelochemicals can definitely reduce the ecological impact, particularly by reducing the amount of herbicide used. Allelopathic rice cultivars combined with cultural management options is, therefore, an interesting and potential technique, contributing to alternative chemical control of weeds in paddy ecosystems (Weston 1996, Olofsdotter 2001, Olofsdotter et al. 2002a). Such an allelopathy-based technique for paddy weed control is the most easily transferable to the low-input management systems prevailing in most Asian rice farming systems (Kong 2008). Therefore, the rice allelopathy may be one of the options in the sustainable weed management strategies.

REFERENCES

- Agrawal, G.K., Rakwal, R., Tamogami, S., Yonekura, M., Kubo, A. and Saji, H. (2002) Chitosan activates defense/stress response(s) in the leaves of *Oryza sativa* seedlings. *Plant Physiology and Biochemistry* 40: 1061-1069.
- Araki, Y. and Kurahashi, Y. (1999) Enhancement of phytoalexin synthesis during rice blast infection of leaves by pre-treatment with carpropamid. *Journal of Pesticide Science* 24; 369-374.
- Azmi, M., Abdullah, M.Z. and Fujii, Y. (2000) Exploratory study on allelopathic effect of selected Malaysian rice varieties and rice field weed species. *Journal of Tropical Agricultural Food Science* 28; 39-54.
- Bais, H.P., Park, S.-W., Weir, T.L., Callaway, R.M. and Vivanco, J.M. (2004) How plants communicate using the underground information superhighway. *Trends in Plant Science* 9; 26-32.
- Belz, R.G. (2007) Allelopathy in crop/weed interactions - an update. *Pest Management Science* 63; 308-326.
- Cartwright, D., Langcake, P., Pryce, R.J., Leworthy, D.P. and Ride, J.P. (1977) Chemical activation of host defence mechanisms as a basis for crop protection. *Nature* 267; 511-513.
- Cartwright, D.W., Langcake, P., Pryce, R.J., Leworthy, D.P. and Ride, J.P. (1981) Isolation and characterization of two phytoalexins from rice as momilactones A and B. *Phytochemistry* 20; 535-537.
- Chung, I.-M., Hahn, S.-J. and Ahmad, A. (2005) Confirmation of potential herbicidal agents in hulls of rice, *Oryza sativa*. *Journal of Chemical Ecology* 31; 1339-1352.
- Dilday, R.H., Nastasi, P. and Smith, R.J. Jr. (1989) Allelopathic observations in rice (*Oryza sativa* L.) to ducksalad (*Heteranthera limosa*). *Proceedings of Arkansas Academic Science* 43; 21-22.
- Dilday, R.H., Lin, J. and Yan, W. (1994) Identification of allelopathy in the USDA-ARS rice germplasm collection. *Australian Journal of Experimental Agriculture* 34; 907-910.
- Dilday, R.H., Yan, W.G., Moldenhauer, K.A.K. and Gravois, K.A. (1998) Allelopathic activity in rice for controlling major aquatic weeds. In: Olofsdotter M. (Ed.) *Allelopathy in Rice*, Manila, Philippines: International Rice Research Institute, pp 7-26.
- Fukuta, M., Xuan, T.D., Deba, F., Tawata, S., Khanh, T.D., and Chung, I.M. (2007) Comparative efficacies in vitro of antibacterial, fungicidal, antioxidant, and herbicidal

activities of momilactones A and B. *Journal of Plant Interaction* 2; 245-251.

Gealy, D.R., Wailes, E.J., Estorninos, L.E. and Chavez, R.S.C. (2003) Rice cultivar differences in suppression of barnyardgrass (*Echinochloa crus-galli*) and economics of reduced propanil rates. *Weed Science* 51; 601-609.

Hawes, M.C., Gunawardena, U., Miyasaka, S. and Zhao, X. (2000) The role of root border cells in plant defense. *Trends in Plant Science* 5; 128-133.

Inderjit (1996) Plant phenolics in allelopathy. *Botanical Review* 62; 186-202.

Inderjit, and Duke, S.O. (2003) Ecophysiological aspects of allelopathy. *Planta* 217; 529-539.

Kato, T., Kabuto, C., Sasaki, N., Tsunagawa, M., Aizawa, H., Fujita, K., Kato, Y. and Kitahara, Y. (1973) Momilactones, growth inhibitors from rice, *Oryza sativa* L. *Tetrahedron Letters* 39; 3861-3864

Kato, T., Tsunakawa, M., Sasaki, N., Aizawa, H., Fujita, K., Kitahara, Y. and Takahashi, N. (1977) Growth and germination inhibitors in rice husks. *Phytochemistry* 16; 45-48.

Kato-Noguchi, H., Ino, T., Sata, N. and Yamamura, S. (2002) Isolation and identification of a potent allelopathic substance in rice root exudates. *Physiologia Plantarum* 115; 401-405.

Kato-Noguchi, H. and Ino, T. (2003a) Rice seedlings release momilactone B into the environment. *Phytochemistry* 63; 551-554.

Kato-Noguchi, H., Ino, T. and Ichii, M. (2003b) Changes in release level of momilactone B into the environment from rice throughout its life cycle. *Functional Plant Biology* 30; 995 - 997.

Kato-Noguchi, H., Ota, K. and Ino, T. (2008a) Release of momilactone A and B from rice plants into the rhizosphere and its bioactivities. *Allelopathy Journal* 22; 321-328.

Kato-Noguchi, H., Ino, T. and Ota, K. (2008b) Secretion of momilactone A from rice roots to the rhizosphere. *Journal of Plant Physiology* 165; 691-696.

Kato-Noguchi, H., Hasegawa, M., Ino, T., Ota, K. and Kujime, H. (2010) Contribution of momilactone A and B to rice allelopathy. *Journal of Plant Physiology* 167; 787-791.

Khanh, T.D., Xuan, T.D., and Chung, I.M. (2007) Rice allelopathy and the possibility for weeds management. *Annals of Applied Biology* 151; 325-339.

Kim, K.U., Shin, D.H., Kim, H.Y., Lee, Z.L. and Olofsdotter, M. (1999) Evaluation of allelopathic potential in rice germplasm. *Korean Journal of Weed Science* 19; 1-9.

Kim, S.Y., Madrid, A.V., Park, S.T., Yang, S.J. and Olofsdotter, M. (2005) Evaluation of rice allelopathy in hydroponics. *Weed Research* 45; 74-79.

Kodama, O., Suzuki, T., Miyakawa, J. and Akatsuka, T. (1988) Ultraviolet-induced accumulation of phytoalexins in rice leaves. *Agricultural and Biological Chemistry* 52; 2469-2473.

Kong, C., Liang, W., Xu, X. and Hu, F. (2004) Release and activity of allelochemicals from allelopathic rice seedlings. *Journal of Agricultural and Food Chemistry* 52; 2861-2865.

- Kong, C. (2008). Rice allelopathy. *Allelopathy Journal* 22, 267-273.
- Lee, C.W., Yoneyama, K., Takeuchi, Y., Konnai, M., Tamogami, S. and Kodama, O. (1999) Momilactones A and B in rice straw harvested at different growth stages. *Bioscience, Biotechnology and Biochemistry* 63; 1318-1320.
- Macías, F.A., Molinillo, J.M.G., Varela, R.M. and Galindo, J.G.G. (2007). Allelopathy –a natural alternative for weed control. *Pest Management Science* 63; 327-348.
- McCully, E. (1999) Roots in soil: unearthing the complexities of roots and their rhizospheres. *Annual Review of Plant Physiology and Plant Molecular Biology* 50; 695-718.
- Nojiri, H., Sugimori, M., Yamane, H., Nishimura, Y., Yamada, A., Shibuya, N., Kodama, O., Murofushi, N. and Omori, T. (1996) Involvement of jasmonic acid in elicitor-induced phytoalexin production in suspension-cultured rice cells. *Plant Physiology* 110; 387-392.
- Olofsdotter, M., Navarez, D., Rebulanan, M. and Streibig, J.C. (1999) Weed-suppressing rice cultivars: Does allelopathy play a role? *Weed Research* 39; 441-454.
- Olofsdotter, M. (2001) Rice: A step toward use of allelopathy. *Agronomy Journal* 93; 3-8.
- Olofsdotter, M., Jensen, L.B. and Courtois, B. (2002a) Improving crop competitive ability using allelopathy: An example from rice. *Plant Breeding* 121; 1-9.
- Olofsdotter, M., Rebulanan, M., Madrid, A., Dali, W., Navarez, D. and Olk, D.C. (2002b) Why phenolic acids are unlikely primary allelochemicals in rice. *Journal of Chemical Ecology* 28; 229-242.
- Putnam, A.R. and Tang, C.-S. (1986) Allelopathy: State of the science. In: Putnam A.R. & Tang C.-S. (Eds.) *The Science of Allelopathy*, New York: John Wiley & Sons, pp 1-19.
- Rice, E.L. (1984) *Allelopathy*, Ed. 2. Orlando: Academic Press:
- Rimando, A.M. and Duke, S.O. (2003) Studies on rice allelochemicals. In: Smith C.W. & Dilday, R.H. (Eds.) *Rice; Origin, History, Technology and Production*, Hoboken, New Jersey: John Wiley & Sons, pp 221-244.
- Seal, A.N., Pratley, J.E., Haig, T. and An, M. (2004a) Identification and quantitation of compounds in a series of allelopathic and non-allelopathic rice root exudates. *Journal of Chemical Ecology* 30; 1647-1662.
- Seal, A.N., Haig, T. and Pratley, J.E. (2004b) Evaluation of putative allelochemicals in rice roots exudates for their role in the suppression of arrowhead root growth. *Journal of Chemical Ecology* 30; 1663-1678.
- Takahashi, N., Kato, T., Tsunagawa, M., Sasaki, N. and Kitahara, Y. (1976) Mechanisms of dormancy in rice seeds. II. New growth inhibitors, momilactone-A and -B isolated from the hulls of rice seeds. *Japanese Journal of Breeding* 26; 91-98.
- Takahashi, A., Kawasaki, T., Henmi, K., Shii, K., Kodama, O., Satoh, H. and Shimamoto, K. (1999) Lesion mimic mutants of rice with alterations in early signaling events of defense. *Plant Journal* 17; 535-545.
- Takeuchi, Y., Kawaguchi, S. and Yoneyama, K. (2001) Inhibitory and promotive allelopathy in rice (*Oryza sativa* L.). *Weed Biology and Management* 1; 147-156.

Tamogami, S. and Kodama, O. (2000) Coronatine elicits phytoalexin production in rice leaves (*Oryza sativa* L.) in the same manner as jasmonic acid. *Phytochemistry* 54; 689-694.

Tesio, F. and Ferrero, A. (2010) Allelopathy, a chance for sustainable weed management. *International Journal of Sustainable Development & World Ecology* 17; 377-389.

Toyomasu, T., Kagahara, T., Okada, K., Koga, J., Hasegawa, M., Mitsuhashi, W., Sassa, T. and Yamane, H. (2008) Diterpene phytoalexins are biosynthesized in and exuded from the roots of rice seedlings. *Bioscience, Biotechnology and Biochemistry* 72; 562-567.

Weston, L.A. (1996) Utilization of allelopathy for weed management in agroecosystems. *Agronomy Journal* 88; 860-866.