

Rice - Golden Apple Snail

Pomacea canaliculata

	<table border="1"> <thead> <tr> <th data-bbox="376 264 887 320">Prevention</th> <th data-bbox="887 264 1200 320">Monitoring</th> <th data-bbox="1200 264 1715 320">Direct Control</th> <th data-bbox="1715 264 1939 320">Direct Control</th> <th data-bbox="1939 264 2163 320">Restrictions</th> </tr> </thead> <tbody> <tr> <td data-bbox="376 320 887 810"> <ul style="list-style-type: none"> • Preferred method: transplanting by 25-30 days old seedling • Prevent snails from entering field by filtering the water entering the field through net traps (5 mm mesh). • Clean equipment (especially harvester) before moving into another field to prevent spreading the pest. • Before planting, make drains in the rice field and add water to the drain. Allow snails to migrate to the water and collect snails and destroy them. • Plant by transplanting rice instead of direct seedling. • Rotate rice with dry land crop and/or allow for fallow periods between flooded rice. </td> <td data-bbox="887 320 1200 810"> <ul style="list-style-type: none"> • Monitor rice seedlings damage by GAS every 2 weeks • Observe egg mass on rice and weeds around rice field once a week • Adults are yellow-brown to greenish brown or dark chestnut, sometimes with dark brown spiral bands of variable number and thickness. 35-60 mm tall • If pest population density exceeds 5 snails per square meter, use Direct Control. </td> <td data-bbox="1200 320 1715 810"> <ul style="list-style-type: none"> • Put sticks (approximately 300 cm long) in rice field. Allow snail to lay eggs and then collect and destroy eggs from the stick. • Drain water from the field and collect the snails with basket/net. • Release ducks into the field (5-10 ducks/ha for 1 month). • If low population, catch by hand and destroy or make snail compost. • Organize field days to collect and destroy snails. • Trap: Take bundles of Papaya, Sweet Potato, Tapioca or Cassava leaves and place them in water along edges of field for at least 24 hours. When the snails get trapped there, collect and destroy them. </td> <td data-bbox="1715 320 1939 395"> <ul style="list-style-type: none"> • Note: for all pesticides usage, follow instructions on product label. </td> <td data-bbox="1939 320 2163 395"></td> </tr> <tr> <td data-bbox="376 395 887 528"> <ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously </td> <td data-bbox="887 395 1200 528"></td> <td data-bbox="1200 395 1715 528"></td> <td data-bbox="1715 395 1939 528"> <ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously </td> <td data-bbox="1939 395 2163 528"></td> </tr> <tr> <td data-bbox="376 528 887 603"> <ul style="list-style-type: none"> • Spray in morning between 6-10 am or in afternoon between 3-6 pm </td> <td data-bbox="887 528 1200 603"></td> <td data-bbox="1200 528 1715 603"></td> <td data-bbox="1715 528 1939 603"> <ul style="list-style-type: none"> • Metaldehyde </td> <td data-bbox="1939 528 2163 603"> <ul style="list-style-type: none"> • WHO Class II (Moderately hazardous) </td> </tr> <tr> <td data-bbox="376 603 887 715"> <ul style="list-style-type: none"> • Niclosamide </td> <td data-bbox="887 603 1200 715"></td> <td data-bbox="1200 603 1715 715"></td> <td data-bbox="1715 603 1939 715"> <ul style="list-style-type: none"> • Copper sulfate </td> <td data-bbox="1939 603 2163 715"> <ul style="list-style-type: none"> • WHO Class IV or U (Unlikely hazardous) • WHO class II </td> </tr> </tbody> </table>	Prevention	Monitoring	Direct Control	Direct Control	Restrictions	<ul style="list-style-type: none"> • Preferred method: transplanting by 25-30 days old seedling • Prevent snails from entering field by filtering the water entering the field through net traps (5 mm mesh). • Clean equipment (especially harvester) before moving into another field to prevent spreading the pest. • Before planting, make drains in the rice field and add water to the drain. Allow snails to migrate to the water and collect snails and destroy them. • Plant by transplanting rice instead of direct seedling. • Rotate rice with dry land crop and/or allow for fallow periods between flooded rice. 	<ul style="list-style-type: none"> • Monitor rice seedlings damage by GAS every 2 weeks • Observe egg mass on rice and weeds around rice field once a week • Adults are yellow-brown to greenish brown or dark chestnut, sometimes with dark brown spiral bands of variable number and thickness. 35-60 mm tall • If pest population density exceeds 5 snails per square meter, use Direct Control. 	<ul style="list-style-type: none"> • Put sticks (approximately 300 cm long) in rice field. Allow snail to lay eggs and then collect and destroy eggs from the stick. • Drain water from the field and collect the snails with basket/net. • Release ducks into the field (5-10 ducks/ha for 1 month). • If low population, catch by hand and destroy or make snail compost. • Organize field days to collect and destroy snails. • Trap: Take bundles of Papaya, Sweet Potato, Tapioca or Cassava leaves and place them in water along edges of field for at least 24 hours. When the snails get trapped there, collect and destroy them. 	<ul style="list-style-type: none"> • Note: for all pesticides usage, follow instructions on product label. 		<ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously 			<ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously 		<ul style="list-style-type: none"> • Spray in morning between 6-10 am or in afternoon between 3-6 pm 			<ul style="list-style-type: none"> • Metaldehyde 	<ul style="list-style-type: none"> • WHO Class II (Moderately hazardous) 	<ul style="list-style-type: none"> • Niclosamide 			<ul style="list-style-type: none"> • Copper sulfate 	<ul style="list-style-type: none"> • WHO Class IV or U (Unlikely hazardous) • WHO class II 		
Prevention		Monitoring	Direct Control	Direct Control	Restrictions																							
<ul style="list-style-type: none"> • Preferred method: transplanting by 25-30 days old seedling • Prevent snails from entering field by filtering the water entering the field through net traps (5 mm mesh). • Clean equipment (especially harvester) before moving into another field to prevent spreading the pest. • Before planting, make drains in the rice field and add water to the drain. Allow snails to migrate to the water and collect snails and destroy them. • Plant by transplanting rice instead of direct seedling. • Rotate rice with dry land crop and/or allow for fallow periods between flooded rice. 		<ul style="list-style-type: none"> • Monitor rice seedlings damage by GAS every 2 weeks • Observe egg mass on rice and weeds around rice field once a week • Adults are yellow-brown to greenish brown or dark chestnut, sometimes with dark brown spiral bands of variable number and thickness. 35-60 mm tall • If pest population density exceeds 5 snails per square meter, use Direct Control. 	<ul style="list-style-type: none"> • Put sticks (approximately 300 cm long) in rice field. Allow snail to lay eggs and then collect and destroy eggs from the stick. • Drain water from the field and collect the snails with basket/net. • Release ducks into the field (5-10 ducks/ha for 1 month). • If low population, catch by hand and destroy or make snail compost. • Organize field days to collect and destroy snails. • Trap: Take bundles of Papaya, Sweet Potato, Tapioca or Cassava leaves and place them in water along edges of field for at least 24 hours. When the snails get trapped there, collect and destroy them. 	<ul style="list-style-type: none"> • Note: for all pesticides usage, follow instructions on product label. 																								
<ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously 			<ul style="list-style-type: none"> • Note: to avoid the development of resistance to pesticides, pesticides in the same FRAC Group should not be used continuously 																									
<ul style="list-style-type: none"> • Spray in morning between 6-10 am or in afternoon between 3-6 pm 			<ul style="list-style-type: none"> • Metaldehyde 	<ul style="list-style-type: none"> • WHO Class II (Moderately hazardous) 																								
<ul style="list-style-type: none"> • Niclosamide 			<ul style="list-style-type: none"> • Copper sulfate 	<ul style="list-style-type: none"> • WHO Class IV or U (Unlikely hazardous) • WHO class II 																								

Cambodia

CREATED/UPDATED: Nov. 2012/ May 2014/Aug 2018

AUTHOR(S): GDA, Phnom Penh, Cambodia

EDITED BY: Plantwise

