



The Prevalence of Virulence Genes and Virulotypes of *Escherichia coli* Strains Isolated from Hospital Wastewaters in Tehran, Iran

***Reza RANJBAR¹, Omid FARAHANI²**

1. Molecular Biology Research Center, Systems Biology and Poisonings Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran
2. Dept. of Microbiology, Islamic Azad University, Varamin-Pishva Branch, Tehran, Iran

***Corresponding Author:** Email: ranjbarre@gmail.com

(Received 16 Jun 2017; accepted 10 Dec 2017)

Abstract

Background: Due to the widespread different pathogenic strains, *Escherichia coli* lead many severe to normal diseases worldwide. Finding the relation of clones with genomic content and clinical features is a key point to recognize the high potential-invasive strains. Specific virulence factors include adhesions, invasions, toxins, and capsule are the main determinants of pathogenic factors of *E. coli* strains.

Methods: From Jun 2014 to Jun 2016, *E. coli* isolates recovered using standard bacteriological methods from wastewater sources in different hospitals in Tehran, Iran, were monitored to recognize the virulence genes by polymerase chain reaction (PCR) assay.

Results: The high and low presences of virulence factors were *fimH*, 76% and *afa*, 13%, respectively.

Conclusion: The results indicated the potential pathogenicity of *E. coli* strains circulating in hospital wastewaters in Tehran, Iran.

Keywords: *Escherichia coli*, Hospital wastewater, Virulence factors

Introduction

As one of the main sources of risk issues related to public health, there is an increasing concern about water contaminations despite recent development in the field of water treatment technologies (1). Coastal zones with the high percentage of untreated sewage injected into waters play a significant role in terms of pollution formation supposed to be the main cause for the greatest number of public health hazard worldwide (2-4) containing waterborne severe diseases and infections like typhoid fever, cholera, dysentery and traveller's diarrhea, caused by different types of bacterial pathogens (5). Expanding the spectrum of waterborne infections (6), lack of regular monitoring of waterborne pathogens, as well as accurate and cost-effective diagnostic tests, are the major issues required to be concerned in order to

protect public health, prevent and control the infections transmitted by waterborne pathogens. The presence of *E. coli* in a variety of infections (7-9), and its diverse clinical spectrum ranging from asymptomatic bacteriuria (reported to have a prevalence of 4%–6% in young women and up to 20% in elderly women (10, 11)) to acute cystitis to pyelonephritis is considered as one of the most common troublesome medicinal issue in an extensive variety of countries all around the world. The complex pathogenesis analyses are infected by both the host biological and behavioral, and the virulence characteristics of the infecting uropathogenic (12). Its propensity to rehash and the certifiable test posed to practical chemotherapy make it worth focus on the virulence repertoire of *E. coli* strains isolated from infected samples.

The range of well-recognized virulence factors (VF), for instance, fimbrial and afimbrial adhesins, toxins and siderophores and new potential ones have been implicated to *E. coli* strains to agitate the host prosperity congruity (13). Many genes are involved, which means that interaction of various virulence genes with host immune system resulted in infection. The exact role of most virulence determinants, however, is still not clearly known in many infection pathogenesis (14).

Conventional methods for detection of *E. coli* based on culturing were time-consuming and not enough efficient in order to identification of pathogenic strains. Improvement of molecular technology has conducted apprehensive studies in the field of uropathogenic *E. coli* (8, 12, 15-24). The convenient and rapid approach of polymerase chain reaction (PCR) leads to figure out an outstanding assessment of virulence determinants detected which may be useful for diagnosis applications and therapeutic strategies (25-28). Unfortunately, despite the importance of uropathogenic *E. coli*, few studies have been performed for monitoring of uropathogenic *E. coli* especially in waters and wastewaters samples (29).

Recent studies are carried out interested to assess the prevalence of different operons coding for virulence factors among *E. coli* strains isolated from various samples from different sources and establish the data bank containing the molecular characteristics of *E. coli* strains to circulate in different countries.

In the present study, *E. coli* isolates from wastewater sources of different hospitals in Tehran, Iran, were

monitored to recognize the virulence genes by polymerase chain reaction (PCR) assay.

Materials and Methods

Sample collection and processing

E. coli strains isolated from hospital wastewater sources over a 24-months sampling period (Jun 2014-Jun 2016) were included in the study. For further analysis, sampling bottles were shipped to the Laboratory in Molecular Biology Research Center, Baqiyatallah, Tehran, Iran. Standard Methods for the Examination of Water and Wastewater 22nd edition were used as the basic methods for culture-based tests (30). All samples were analyzed for the presence of *E. coli*. *E. coli* colonies from agar plates were picked and streaked for purity on EMB agar. Well-isolated colonies of purified *E. coli* were stored for long-term storage in SKIM Milk media with 20% (v/v) glycerol and at -20 °C. The *E. coli* strains were isolated by minimal standard bacteriological test and identified compared with the *E. coli* laboratory reference using conventional biochemical tests.

PCR

E. coli isolates were tested for the presence of specific primers used to amplify sequences of the type 1 fimbriae (*fimH*), pili associated with pyelonephritis (*pap*), S fimbriae (*sfa*), afimbrial adhesins (*afa*), hemolysin (*hly*), and cytotoxic necrotizing factor (*cnf*), aerobactin (*aer*). Details of primer sequences, predicted sizes of the amplified products and specific annealing temperatures are given in Table 1.

Table 1: Characteristics of primer sequences used for PCR assays

Virulence factor	Target gene(s)	Primer sequence (5'-3')	Size fragment (bp)
Type 1 fimbriae	<i>fimH</i>	F:GAGAAGAGGTTTGATTAACTTATTG R:AGAGCCGCTGTAGAAGCTGAGG	559
P fimbriae	<i>pap</i>	F:GACGGCTGTACTGCAGGG TGTGGCG R:ATATCCTTTCTGCAGGGATGCAATA	328
Afa adhesins	<i>afa</i>	F:GCTGGGCAGCAAAGCTGATAAC TCTC R:CATCAAGCTGTTTGTTCGTCGGCCG	750
Haemolysin	<i>hlyA</i>	F:AACAAGGATAAGCACTGTTCTGGCT R:ACCATATAAGCGGTCAATCCCCGTCA	1177
Cytotoxic necrotizing factor	<i>cnf</i>	F:TTATATAGTCGTCAAGATGGA R:CACTAAGCTTTACAATATTGA	639
Aerobactin	<i>aer</i>	F:TACCGGATTGTTCATATGCAGACCGT R:AATATCTTCCTCCAGTCCGGAGAAG	602

Detection of sequences of target genes was done by PCR (31). PCR amplification of bacterial DNA extracts was performed in 20 µl of template DNA (1 µl), 10 pmoles of each primer, the four deoxynucleoside triphosphates (each at 10 µM) and 1.25 U Taq DNA polymerase in 2x PCR buffer containing MgCl₂. For finding the optimum amplification procedure, the denaturation and annealing conditions were described in Table 2. The final reaction mixture (6µl) underwent gel electrophoresis in 1.5% agarose containing in 100 ml TBE (1X) buffer (1,6 M Tris-EDTA, 0,02 M boric acid) per

ml. Amplified DNA fragments of specific sizes were detected by UV induced fluorescence. Overall, 1000 bp DNA ladder (Viogene) was used for determining the sizes of the amplicons.

The amplified DNA product was visualized by standard submarine gel electrophoresis using 10 µL of the final reaction mixture on a 2% agarose gel. Amplified DNA fragments of specific sizes were located by UV fluorescence, after staining with ethidium bromide. The 100-bp ladder was used as a standard for determining molecular size of PCR products (Fig. 1).

Table 2: PCR conditions of target genes

<i>Genes</i>	<i>Initial denaturation</i> (°C/min)	<i>Denaturation</i> (°C/s)	<i>Annealing</i> (°C/s)	<i>Extension</i> (°C/s)	<i>Final extension</i> (°C/min)
<i>fimH</i>	95/5	94/60	59/60	72/60	72/7
<i>pap</i>	95/5	94/60	63/60	72/60	72/7
<i>afa</i>	95/5	94/60	65/60	72/60	72/7
<i>blyA</i>	95/5	94/60	62/60	72/60	72/7
<i>cnf</i>	95/5	94/60	57/60	72/60	72/7
<i>aer</i>	95/5	94/60	68/60	72/60	72/7




Fig. 1: Detection of *fimH* (559 bp), *aer* (602 bp), *pap* (328 bp), and *afa* (750 bp) gene by PCR in hospital wastewaters. Lane 5 and 6: negative representative *blyA* and *cnf* genes

Results

Eighty strains of *E. coli* were isolated from hospital wastewater sources and subjected to PCR as-

say in order to determine the prevalence of virulence genes.

The frequencies and profiles of virulence genes are shown in Table 3 and 4, respectively. The prevalence of genes of fimbrial adhesive systems was 76% and 16% for *fimH*, and *pap*, respectively. *Afa* afimbrial adhesins were detected in 13% of strains. The prevalence of 53% was identified for the *aer* gene. The presence of two genes encoding toxins, i.e. *bly* and *cnf* were not detected. Sixty-one strains shared DNA sequences related to *fimH* and other genes associated as *aer*, *pap*, and *afa*. *FimH* adhesin was the most prevalent virulence factor detected, having occurred in 76% of strains, and it was detected in 16% of operon *pap*. The studied strains of the total *fimH* showed adhesin-encoding operon *fimH* (95%), *fimH-afa* (20%), *fimH-afa-pap* (2.5%) and *fimH-pap* (1.2%) and 51 of them showed the other VF gene: *Aer-fimH* (47%), *Aer-fimH-afa* (6.2%) and *Aer-fimH-pap* (10%) (Table 4). The high presence of adhesion genes and the low presence of genes encoding toxins were illustrated in our finding.

Table 3: Frequency of virulence genes among 80 *E. coli* isolates

<i>Virulence gene</i>	<i>Total (n = 80), n (%)</i>
<i>fimH</i>	61(76)
<i>aer</i>	43(53)
<i>Pap</i>	13(16)
<i>afa</i>	11(13)
<i>blyA</i>	0 (0)
<i>cnf</i>	0 (0)

Table 4: Nine different patterns for virulence gene profiles

<i>Profile names</i>	<i>Virulence gene patterns</i>	<i>Total (n = 80), n (%)</i>
[Type 1]	<i>fimH-afa</i>	16 (20)
[Type 2]	<i>fimH-afa-pap</i>	2 (2.5)
[Type 3]	<i>afa-pap</i>	2(2.5)
[Type 4]	<i>fimH-pap</i>	1(1.2)
[Type 5]	<i>Aer- fimH</i>	38(47)
[Type 6]	<i>Aer- fimH-afa</i>	5(6.2)
[Type 7]	<i>Aer- fimH-pap</i>	8(10)
[Type 8]	<i>Aer- afa</i>	5(6.2)
[Type 9]	<i>Aer-pap</i>	8(10)

Discussion

Finding the relation of genetic virulence traits of the clinical isolates and the type of infections will influence the treatment of these infections. Adhesins, toxins and iron-chelating factors (siderophores) were reported as the main virulence factors (VFs) of *E. coli* (32) while afimbrial adhesins (*afa*), pfimbrial adhesins (*pap*) and s-fimbrial adhesins (*sfa*) were indicated as those adhesins by which UPEC typically attach to epithelial cells lining (33).

Intimin, as one of the pathogenic factor, was playing an important role in virulence helping bacteria to attach the epithelial cell lining (28, 34). A recent research about cytotoxic necrotizing and haemolytic factors was conducted (18). Forty percent of UPEC produce cytotoxic necrotizing factor 1(*cnf1*) involved in the dissemination and persistence of cells in the urinary tract. Aerobactin, the iron chelating agent, encoded by *aer* have been reported as a determinate factor to enabling colonization of *E. coli* in iron-poor environments (35). Promotion of bacterial invasion through the

epithelial barrier is considered due to the cytolytic effect of α -hemolysin (*blyA*) encoded by *a-bly*. In this study, the high percentage of adhesins gene could be related to the pathogenicity of the isolated strains as adherence is the most important pathogenicity determinant (36). *fimH* was highly conserved in UTI isolates, which confirms its crucial role during colonization of the urinary tract (36-39). The presence of *pap* gene had been considered in association with pyelonephritis; therefore, lower percentages of *pap* gene (16%) may propose the lower capabilities of isolated strains to colonize kidneys and generate pyelonephritis (40, 41). The *blyA* and *cnf* genes indicated the positive relationship with pathogenicity island PAI II₉₆, and the pathogenicity island PAI I_{CF1073} is related to *aer* gene as well as with *blyA* and *pap* operon (42-46).

Molecular methods particularly DNA-based detection techniques help us help us in the field of diagnostic and molecular epidemiology of pathogenic microbial strains (47-48). The technique used in current study was able of to elucidate

successfully the pathogenicity status of the strains under the study.

Conclusion

Consequently, we represent the high frequency of *fimH* gene among isolated strains of *E. coli* from wastewater samples followed by new patterns of frequencies in association with virulence genes of *aer*, *pap*, *afa*, and *fimH*. PCR typing acts as a useful method to determine the virulence gene profiles of *E. coli* strains and detect the complete data from genome content in association with different *E. coli* strains that may cause severe illnesses worldwide. The high percent of virulence genes of *E. coli* strains circulating in hospital wastewaters indicated a warning danger for the transfer of these genes to the environment. Therefore, immediate measures are needed to control and treat hospital wastewaters more effectively.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been considered by the authors.

Acknowledgments

We would like to thank from the “Clinical Research Development Center of Baqiyatallah hospital” for their kind cooperation. This study was financially supported in part by “Clinical Research Development Center of Baqiyatallah hospital.

Conflict of interest

The authors declare that there is no conflict of interests.

References

1. Kümmerer K (2004). Resistance in the environment. *J Antimicrob Chemother*, 54(2): 311-20.
2. Lee D-Y, Shannon K, Beaudette LA (2006). Detection of bacterial pathogens in municipal wastewater using an oligonucleotide microarray and real-time quantitative PCR. *J Microbiol Methods*, 65(3): 453-67.
3. Ranjbar R, Pezeshknejad P, Khamesipour F et al (2017). Genomic fingerprints of *Escherichia coli* strains isolated from surface water in Alborz province, Iran. *BMC Res Notes*, 10 :295.
4. Ranjbar R, Khamesipour F, Hosseini S (2017). Prevalence and distribution of *E. coli* and its virulence genes from different water sources in Alborz province, Iran. *J Biochem Tech*, 7(2): 1116-22.
5. Hunter P (2003). Climate change and waterborne and vector-borne disease. *J Appl Microbiol*, 94 Suppl:37S-46S.
6. Marshall MM, Naumovitz D, Ortega Y, Sterling CR (1997). Waterborne protozoan pathogens. *Clin Microbiol Rev*, 10(1): 67-85.
7. Ronald A (2002). The etiology of urinary tract infection: traditional and emerging pathogens. *Dis Mon*, 49(2):71-82.
8. Ranjbar R, Masoudimanesh M, Dehkordi FS et al (2017). Shiga (Vero)-toxin producing *Escherichia coli* isolated from the hospital foods; virulence factors, o-serogroups and antimicrobial resistance properties. *Antimicrob Resist Infect Control*, 6: 4.
9. Ranjbar R, Sheikhshahrokh A, Jonaidi Jafari N (2017). Shiga (vero) toxin producing *Escherichia coli* in various types of food stuffs; virulence factors, O- serogroups and antimicrobial resistance properties. *J Food Saf*, 37: e12312.
10. Lindberg U, Claesson I, Hanson LÅ, Jodal U (1978). Asymptomatic bacteriuria in schoolgirls: VIII. Clinical course during a 3-year follow-up. *J Pediatr*, 92(2): 194-9.
11. Hooton TM, Scholes D, Stapleton AE et al (2000). A prospective study of asymptomatic bacteriuria in sexually active young women. *N Engl J Med*, 343(14): 992-7.
12. Usein CR, Damian M, Tatu- Chitoiu D et al (2001). Prevalence of virulence genes in *Escherichia coli* strains isolated from Romanian adult urinary tract infection cases. *J Cell Mol Med*, 5(3): 303-10.
13. Stevens DL, Bisno AL, Chambers HF et al (2005). Practice guidelines for the diagnosis

- and management of skin and soft-tissue infections. *Clin Infect Dis*, 41(10):1373-406.
14. Agarwal J, Srivastava S, Singh M (2012). Pathogenomics of uropathogenic *Escherichia coli*. *Indian J Med Microbiol*, 30(2): 141-9.
 15. Zhang L, Foxman B, Manning SD et al (2000). Molecular epidemiologic approaches to urinary tract infection gene discovery in uropathogenic *Escherichia coli*. *Infect Immun*, 68(4): 2009-15.
 16. Johnson JR, Russo TA, Tarr PI et al (2000). Molecular epidemiological and phylogenetic associations of two novel putative virulence genes, *iha* and *iroN*(*E. coli*), among *Escherichia coli* isolates from patients with urosepsis. *Infect Immun*, 68(5): 3040-7.
 17. Momtaz H, Karimian A, Madani M et al (2013). Uropathogenic *Escherichia coli* in Iran: serogroup distributions, virulence factors and antimicrobial resistance properties. *Ann Clin Microbiol Antimicrob*, 12: 8.
 18. Tajbakhsh E, Khamesipour F, Ranjbar R, Ugwu IC (2015). Prevalence of class 1 and 2 integrons in multi-drug resistant *Escherichia coli* isolated from aquaculture water in Chaharmahal Va Bakhtiari province, Iran. *Ann Clin Microbiol Antimicrob*, 14: 37.
 19. Farshad S, Ranjbar R, Japoni A et al (2012). Microbial susceptibility, virulence factors, and plasmid profiles of uropathogenic *Escherichia coli* strains isolated from children in Jahrom, Iran. *Arch Iran Med*, 15(5): 312-316.
 20. Jahandeh N, Ranjbar R, Behzadi P, Behzadi E (2015). Uropathogenic *Escherichia coli* virulence genes: invaluable approaches for designing DNA microarray probes. *Cent European J Urol*, 68(4): 452-8.
 21. Anvarinejad M, Farshad S, Ranjbar R et al (2012). Genotypic analysis of *E. coli* strains isolated from patients with cystitis and pyelonephritis. *Iran Red Crescent Med J*, 14(7): 408-416.
 22. Ranjbar R, Haghi-Ashtiani M, Jafari NJ, Abedini M (2009). The prevalence and antimicrobial susceptibility of bacterial uropathogens isolated from pediatric patients. *Iran J Public Health*, 38(2): 134-8.
 23. Ranjbar R, Hosseini S, Zahraei-Salehi T et al (2016). Investigation on prevalence of *Escherichia coli* strains carrying virulence genes *ipaH*, *estA*, *eaeA* and *bfpA* isolated from different water sources. *Asian Pac J Trop Dis*, 6(4): 278-83.
 24. Khademestarki NS, Ranjbar R (2016). The phylogenetic study of *Escherichia coli* strains isolated from clinical cases. *J Pure Appl Microbiol*, 10(1): 351-5.
 25. Johnson JR, Stell AL (2000). Extended virulence genotypes of *Escherichia coli* strains from patients with urosepsis in relation to phylogeny and host compromise. *J Infect Dis*, 181(1): 261-72.
 26. Foxman B, Zhang L, Tallman P et al (1995). Virulence characteristics of *Escherichia coli* causing first urinary tract infection predict risk of second infection. *J Infect Dis*, 172(6): 1536-41.
 27. Johnson JR, Brown JJ, Carlino UB, Russo TA (1998). Colonization with and acquisition of uropathogenic *Escherichia coli* as revealed by polymerase chain reaction-based detection. *J Infect Dis*, 177(4): 1120-4.
 28. Le Bouguenec C, Archambaud M, Labigne A (1992). Rapid and specific detection of the *pap*, *afa*, and *sfa* adhesin-encoding operons in uropathogenic *Escherichia coli* strains by polymerase chain reaction. *J Clin Microbiol*, 30(5):1189-93.
 29. Ranjbar R, Farahani O (2017). The Prevalence of Plasmid-mediated Quinolone Resistance Genes in *Escherichia coli* Isolated from Hospital Wastewater Sources in Tehran, Iran. *Iran J Public Health*, 46(9): 1285-91.
 30. APHA/AWWA/WEF (2012). *Standard methods for the examination of water and wastewater*. 22nd edn American public health Association/American water works Association/Water environment federation, Washington DC, USA
 31. Rendón MaA, Saldaña Z, Erdem AL et al (2007). Commensal and pathogenic *Escherichia coli* use a common pilus adherence factor for epithelial cell colonization. *Proc Natl Acad Sci U S A*, 104(25): 10637-42.
 32. Jauréguy F, Carbonnelle E, Bonacorsi S et al (2007). Host and bacterial determinants of initial severity and outcome of *Escherichia coli* sepsis. *Clin Microbiol Infect*, 13(9): 854-62.
 33. Dobrindt U, Blum-Oehler G, Hartsch T et al (2001). S-Fimbria-Encoding Determinant *sfaI* Is Located on Pathogenicity Island III536 of

- Uropathogenic *Escherichia coli* Strain 536. *Infect Immun*, 69(7): 4248-56.
34. Le Bouguenec C, Garcia M-I, Ouin V et al (1993). Characterization of plasmid-borne afa-3 gene clusters encoding afimbrial adhesins expressed by *Escherichia coli* strains associated with intestinal or urinary tract infections. *Infect Immun*, 61(12): 5106-14.
 35. Wiles TJ, Kulesus RR, Mulvey MA (2008). Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol*, 85(1): 11-9.
 36. Mulvey MA (2002). Adhesion and entry of uropathogenic *Escherichia coli*. *Cell Microbiol*, 4(5): 257-71.
 37. Guyer DM, Gunther IV NW, Mobley HL (2001). Secreted proteins and other features specific to uropathogenic *Escherichia coli*. *J Infect Dis*, 183 Suppl 1:S32-5.
 38. Bower JM, Eto DS, Mulvey MA (2005). Covert operations of uropathogenic *Escherichia coli* within the urinary tract. *Traffic*, 6(1): 18-31.
 39. Antão E-M, Wieler LH, Ewers C (2009). Adhesive threads of extraintestinal pathogenic *Escherichia coli*. *Gut Pathog*, 1(1): 22.
 40. Lane M, Mobley H (2007). Role of P-fimbrial-mediated adherence in pyelonephritis and persistence of uropathogenic *Escherichia coli* (UPEC) in the mammalian kidney. *Kidney Int*, 72(1): 19-25.
 41. Yamamoto S (2007). Molecular epidemiology of uropathogenic *Escherichia coli*. *J Infect Chemother*, 13(2): 68-73.
 42. Bingen-Bidois M, Clermont O, Bonacorsi S et al (2002). Phylogenetic analysis and prevalence of urosepsis strains of *Escherichia coli* bearing pathogenicity island-like domains. *Infect Immun*, 70(6): 3216-26.
 43. Soto S, Zuniga S, Ulleryd P, Vila J (2011). Acquisition of a pathogenicity island in an *Escherichia coli* clinical isolate causing febrile urinary tract infection. *Eur J Clin Microbiol Infect Dis*, 30(12): 1543-50.
 44. Hancock V, Ferrieres L, Klemm P (2008). The ferric yersiniabactin uptake receptor FyuA is required for efficient biofilm formation by urinary tract infectious *Escherichia coli* in human urine. *Microbiology*, 154(Pt 1): 167-75.
 45. Clarke TE, Tari LW, Vogel HJ (2001). Structural biology of bacterial iron uptake systems. *Curr Top Med Chem*, 1(1): 7-30.
 46. Rowe MC, Withers HL, Swift S (2010). Uropathogenic *Escherichia coli* forms biofilm aggregates under iron restriction that disperse upon the supply of iron. *FEMS Microbiol Lett*, 307(1): 102-9.
 47. Ranjbar R, Karami A, Farshad S, Giammanco GM, Mammina C (2014). Typing methods used in the molecular epidemiology of microbial pathogens: a how-to guide. *New Microbiol*, 37(1):1-15.
 48. Khakabimamaghani S, Najafi A, Ranjbar R, Raam M (2013). GelClust: a software tool for gel electrophoresis images analysis and dendrogram generation. *Comput Methods Programs Biomed*, 111 (2):512-8.