




Green Area in Urban Cuiabá: The Case of the Universidade Federal do Mato Grosso

Poliane Pierra Patricio Pazini¹  0000-0002-5924-809X

Diego Tyszka Martinez²  0000-0002-7275-8043

Sidney Fernando Caldeira¹  0000-0001-6042-4313

Rodrigo Adversi Silva¹  0000-0002-5849-0386

Felipe Andrade Antonieto³  0000-0001-8844-5140

Abstract

Urban trees/vegetation are one of the factors related to a better quality of life of the population. Therefore, knowledge on the structure and quality of urban green areas is important to implement actions that intensify this benefit. This study aimed to quantify the floristic composition of the Universidade Federal do Mato Grosso, in Cuiabá. A census was conducted to count individuals with soil circumference ≥ 15.7 cm, wherein 190 species were identified, with Fabaceae being the most representative family at 24.2%. Of the total, 23.7% are exotic species to Brazil and 62.6% are native to the Cerrado. The most abundant species were *Licania tomentosa*, with 8.9% of individuals, *Mangifera indica* at 7.8%, *Jacaranda cuspidifolia* with 6.3%, *Handroanthus impetiginosus* with 5.4%, and *Anadenanthera peregrine* with 4.4%. The university campus houses a high floristic richness and diversity of species, with an adequate proportion of Brazilian native species. However, its composition needs to be adapted to comply with municipal legislation.

Keywords: afforestation, diversity, public spaces.

1. INTRODUCTION AND OBJECTIVES

Human occupation in cities most often increases fast, in disorder and without planning, with direct consequences on quality of life (Pinheiro & Souza, 2017). In this way, performing and deepening studies related to urban afforestation is necessary, due to the contribution of urban greenery to an improved environmental quality; a fact that also generates positive psychological and physical effects for its inhabitants (McHale et al., 2007).

Green areas are the results of landscaping processes in cities which have several influence of man, and which divide cultural green areas such as plazas, squares and gardens, and urban forest fragments, which are basically anthropized forest remnants due to the urbanization process (Biondi, 2015).

The ecological function of these areas is related to the quality of the urban environment, favoring climate mitigation since the natural elements that compose these spaces attenuate the

impacts resulting from industrialization. There is a feeling of well-being in these places in different environments, which enables community permanence and also provides areas for meetings and social relationships (Loboda & Angelis, 2005).

Green areas structure ecological corridors, especially with the presence of native fruit species, which helps developing a richer fauna and is a very important element for maintaining the biodiversity of urban ecosystems and indicators of environmental quality in cities (Alencar & Cardoso, 2015).

Knowledge of the floristic composition of urban green areas is fundamental for environmental balance and should be part of frequent study programs in order to value the landscape and the local physiognomy with implantation of species adapted to each situation. In addition to the benefits of vegetation to human health, the ecological peculiarity of the region should also be considered in order to preserve the genetic identity (Kramer & Krupek, 2012).

¹ Universidade Federal de Mato Grosso (UFMT), Cuiabá, MT, Brasil

² Universidade Federal do Paraná (UFPR), Curitiba, PR, Brasil

³ Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, AM, Brasil

For urban afforestation to perform its intended function, it is necessary to prior plan planted green areas, since many of the benefits that the population receives from ecosystems in order to guarantee their well-being depends on tree species diversification (Morgenroth et al., 2016). In this sense, Brazilian academic institutions, with their potential for scientific production, should develop floristic projects that consider the peculiar characteristics of the species and use them for implanting forestry with superior quality (Eisenlohr et al., 2008).

The Universidade Federal do Mato Grosso (UFMT) campus is characterized as an important green area fragment, which contributes to the quality of life of local inhabitants and provides environmental services to the city. With an apparent tree flora diversity comprising native and exotic species, it is located in an original Cerrado region, with the most part being artificial trees implanted in its construction phase.

Knowledge of the existing tree species on the campus is important for planning urban afforestation in order to better use the characteristics of each individual in the environment, as well as to know its richness and diversity. This fact supports new native species plantations, and the removal of exotic species if necessary.

In view of the above, this study aimed to learn about the floristic composition of UFMT afforestation at the Cuiabá campus to show the specific aspects that can be implemented to improve its structure.

2. MATERIALS AND METHODS

This study was conducted at the UFMT, Cuiabá campus, circumscribed to the coordinates 15° 35' 46" South latitude and 56° 5' 48" West longitude. The campus extends to an area of about 76 ha in the Boa Esperança neighborhood of Cuiabá, Mato Grosso.

Most of the vegetation that makes up UFMT was planted in the early 1970s, with an apparent tree species diversity native to the Cerrado. Later plantings were carried out over the years by student and commemorative activities or to obtain shading in order to mitigate the effect of heat.

The climate of the region is Aw (Köppen-Geiger) with well-defined dry winter and rainy summer seasons, and average annual temperature of 25 °C. The average local altitude is 150 m, with average annual rainfall ranging between 1,300 mm.year⁻¹ and 1600 mm.year⁻¹ (Alvares et al., 2013).

According to the Embrapa soil classification (Santos et al., 2013), the predominant soil of the region is Dystrophic Red-yellow Argisol. The local vegetation comprises a large diversity

of native and exotic tree species, with emphasis on Cerrado species, a predominant forest typology in the capital of Mato Grosso. The city of Cuiabá has an approximate population of 590,000 inhabitants in an area of 3,293,537 km² (IBGE, 2017).

Data were collected via a census of tree vegetation at UFMT, Cuiabá campus, conducted from July 2016 to January 2017. Data were recorded in pre-structured tables containing tree number and species information. Using a measuring tape, individuals with a circumference at 1.30 m above ground level (breast-height) (CBH 1.3 m) equal to or greater than 15.7 cm were included, which is equivalent to 5 tape cm in diameter at 1.3 m (DBH 1.3 m). Each tree received a metal plate with an identification number, which enabled indexing information on the family, genus, and species when possible.

The known species were identified on site, while the unknown species had their botanical material collected for later identification at the Laboratory of Dendrology, School of Forestry Engineering, UFMT.

The species were organized according to families registered in the Angiosperm Phylogeny Group III (Stevens, 2016), and scientific names were checked on the Missouri Botanical Garden (2016) portal. The species were also classified according to their origin as native or exotic to Brazil, as to their natural occurrence in the phytogeographical domain of the Cerrado to which the municipality of Cuiabá belongs, and as to leaf deciduousness, followed by the reference of the Brazil 2020 Flora program currently under construction (Coppetec, c2019).

After data collection, the species were divided into 4 groups according to their relevant characteristic for use in urban afforestation. The created groups were: Palm trees, which gathered the species of the Arecaceae family; Fruit trees, which grouped species that produce fleshy fruits; Ornamental, with species that show intense and striking flowering; and Others, which gathered the species that did not fit into the previous groups, but provided other environmental regulation services to the city, such as shade, climatic mitigation, air pollution retention, noise retention, atmospheric carbon retention and rainwater interception.

Data were analyzed by descriptive statistics starting from structuring the table, with the calculations of relative abundance by species (Equation 1) being performed through the MS Excel 2016:

$$Ab.R = Ab.A * 100 / N \quad (1)$$

Ab.R: number of individuals of a given species; Ab.A: absolute abundance; N: total number of individuals.

Diversity was determined using the Shannon-Weiner Indexes (Equation 2) and Simpson Index (Equation 3) shown by Odum (2012), calculated by the R program (R Development Core Team, 2015):

$$H' = - \sum ((ni/N) * \ln(ni/N)) \quad (2)$$

H': Shannon Index; ni: number of individuals in the i^{th} species; N: total number of individuals.

$$D = \sum (ni * (ni - 1) / (N(N - 1))) \quad (3)$$

D: Simpson Index; ni: number of individuals in the i^{th} species; N: total number of individuals.

3. RESULTS AND DISCUSSION

We surveyed 4817 shrub and tree individuals, of which two were not classified, while 14 were classified regarding family, 66 were classified in genus and 80 were dead. The richness was 189 forest species, with 182 trees and 7 palm trees, distributed across 136 genera and 48 botanical families.

The most representative family of species was Fabaceae, with 24.2%, followed by Malvaceae (7.9%), Myrtaceae (6.8%), Bignoniaceae (5.8%), and others with values less than 4%. The ten most representative families characterized 63.2% of the total trees. Among the species, *Licania tomentosa* was the most abundant with 8.9%, followed by *Mangifera indica* (7.8%), *Jacaranda cuspidifolia* (6.3%), *Handroanthus impetiginosus* (5.4%), and *Anadenanthera peregrina* (4.4%) (Table 1).

Table 1. Floristic composition, origin (Or.), deciduity (Dec.) groups and relative abundance (RAb.) of species in the afforestation of UFMT campus Cuiabá, 2017.

N	Species, genus, family or absence of classification	Or.	Dec.	Groups				RAb.
				P	F	O	Oth.	
1	<i>Licania tomentosa</i> (Benth.) Fritsch	N	PR				x	8.9
2	<i>Mangifera indica</i> L.	E	PR		x			7.8
3	<i>Jacaranda cuspidifolia</i> Mart.	N*	D			x		6.3
4	<i>Handroanthus impetiginosus</i> (Mart. ex DC.) Mattos	N*	D				x	5.4
5	<i>Anadenanthera peregrina</i> (L.) Speg.	N*	D				x	4.4
6	<i>Cariniana rubra</i> Gardner ex Miers	N*	D				x	3.5
7	<i>Genipa americana</i> L.	N*	D		x			3.5
8	<i>Astronium fraxinifolium</i> Schott	N*	D				x	3.1
9	<i>Acrocomia aculeata</i> (Jacq.) Lodd. ex Mart.	N*	PR	x				3
10	<i>Dipteryx alata</i> Vogel	N*	PR		x			2.8
11	<i>Swietenia macrophylla</i> King	N*	D				x	2.2
12	<i>Cordia glabrata</i> (Mart.) A. DC.	N*	D			x		2.1
13	<i>Psidium guajava</i> L.	E	PR		x			1.6
14	<i>Anacardium occidentale</i> L.	N*	D		x			1.5
15	<i>Hevea brasiliensis</i> (Willd. ex A. Juss.) Müll. Arg.	N	D				x	1.5
16	<i>Cecropia pachystachya</i> Trec.	N*	D				x	1.4
17	<i>Curatella americana</i> L.	N*	D				x	1.3
18	<i>Albizia lebeck</i> (L.) Benth.	E	D				x	1.2
19	<i>Samanea tubulosa</i> (Benth.) Barneby & J.W. Grimes	N*	D		x			1.2
20	<i>Handroanthus avellanadae</i> (Lorentz ex Griseb.) Mattos	N	D			x		1.2
21	<i>Delonix regia</i> (Bojer ex Hook.) Raf.	E	D			x		1.1
22	<i>Myracrodruon urundeuva</i> Allemão	N*	D				x	1.1
23	<i>Inga laurina</i> (Sw.) Willd.	N*	PR				x	1.1
24	<i>Handroanthus aureus</i> Mattos	N*	D			x		1.0
25	<i>Enterolobium contortisiliquum</i> (Vell.) Morong	N*	D				x	1.0
26	<i>Guazuma ulmifolia</i> Pers.	N*	D		x			1.0
27	<i>Vochysia divergens</i> Pohl	N*	PR				x	0.9

Table 1. Continued...

N	Species, genus, family or absence of classification	Or.	Dec.	Groups				RAb.
				P	F	O	Oth.	
28	<i>Mimosa caesalpiniiifolia</i> Benth.	N*	D				x	0.9
29	<i>Cassia siamea</i> Lam.	N*	PR				x	0.8
30	<i>Gmelina arborea</i> Roxb. ex Sm.	E	D		x			0.7
31	<i>Anadenanthera falcata</i> (Benth.) Speg.	N*	D				x	0.7
32	<i>Handroanthus serratifolius</i> (Vahl) S.O. Grose	N*	D				x	0.7
33	<i>Leucaena leucocephala</i> (Lam.) de Wit	E	D				x	0.6
34	<i>Cordia aff. myxa</i> L.	E	PR				x	0.6
35	<i>Cocos nucifera</i> L.	N	PR	x				0.6
36	<i>Caryocar brasiliense</i> A. St.-Hil.	N*	D		x			0.6
37	<i>Artocarpus heterophyllus</i> Lam.	N	PR		x			0.5
38	<i>Buchenavia tomentosa</i> Eichler	N*	D		x			0.5
39	<i>Casearia sylvestris</i> Sw.	N*	D		x			0.5
40	<i>Magonia pubescens</i> A. St.-Hil.	N*	D				x	0.5
41	<i>Pithecellobium hassleri</i> Chodat	N	D				x	0.5
42	<i>Vatairea macrocarpa</i> (Benth.) Ducke	N*	D				x	0.5
43	<i>Tamarindus indica</i> L.	E	D		x			0.5
44	<i>Bowdichia virgilioides</i> Kunth	N*	D				x	0.4
45	<i>Amburana acreana</i> (Ducke) A.C. Sm.	N	D				x	0.4
46	<i>Ceiba boliviana</i> Britten & Baker f.	N*	D				x	0.4
47	<i>Talisia esculenta</i> (A. St.-Hil.) Radlk.	N*	D		x			0.4
48	<i>Handroanthus heptaphyllus</i> (Vell.) Mattos	N*	D				x	0.4
49	<i>Simarouba versicolor</i> St.-Hil.	N*	D		x			0.4
50	<i>Hymenaea courbaril</i> L.	N*	D		x			0.3
51	<i>Luehea paniculata</i> Mart.	N*	D				x	0.3
52	<i>Qualea grandiflora</i> Mart.	N*	D				x	0.3
53	<i>Terminalia catappa</i> L.	E	D				x	0.3
54	<i>Ficus benamina</i> L.	N*	PR		x			0.3
55	<i>Inga</i> sp.	N*	PR				x	0.3
56	<i>Protium heptaphyllum</i> (Aubl.) Marchand	N*	PR		x			0.3
57	<i>Cassia fistula</i> L.	E	D				x	0.3
58	<i>Hymenaea stigonocarpa</i> Mart. ex Hayne	N*	D		x			0.3
59	<i>Vochysia haenkeana</i> Mart.	N*	D				x	0.3
60	<i>Aspidosperma riedelii</i> Müll.Arg	N	D				x	0.3
61	<i>Erythroxylum suberosum</i> A. St.-Hil.	N*	D		x			0.3
62	<i>Eucalyptus</i> sp. 2	E	D				x	0.3
63	<i>Luehea grandiflora</i> Mart.	N*	D				x	0.3
64	<i>Murraya paniculata</i> (L.) Jack	E	PR		x			0.3
65	<i>Peltophorum dubium</i> (Spreng.) Taub.	N*	D				x	0.3
66	<i>Spondias purpurea</i> L.	E	D		x			0.3
67	<i>Cedrela fissilis</i> Vell.	N*	D				x	0.2
68	<i>Chorisia speciosa</i> A. St.-Hil.	N*	D				x	0.2
69	<i>Cordia trichotoma</i> (Vell.) Arráb. ex Steud.	N*	D				x	0.2
70	<i>Eucalyptus</i> sp.	E	D				x	0.2

Table 1. Continued...

N	Species, genus, family or absence of classification	Or.	Dec.	Groups				RAb.
				P	F	O	Oth.	
71	<i>Handroanthus roseo-albus</i> (Ridl.) Mattos	N*	D			x		0.2
72	<i>Physocalymma scaberrimum</i> Pohl	N*	D			x		0.2
73	<i>Platypodium elegans</i> Vogel	N*	D			x		0.2
74	<i>Andira cuyabensis</i> Benth.	N*	D				x	0.2
75	<i>Clitoria fairchildiana</i> R.A. Howard	N*	D				x	0.2
76	<i>Copaifera langsdorffii</i> Desf.	N*	D				x	0.2
77	<i>Malpighia glabra</i> L.	E	PR		x			0.2
78	<i>Muntingia calabura</i> L.	N	D		x			0.2
79	<i>Pachira aquatica</i> Aubl.	N	D		x			0.2
80	<i>Terminalia argentea</i> Mart. ex Succ.	N*	D				x	0.2
81	<i>Guettarda viburnoides</i> Cham. & Schltld.	N*	D		x			0.2
82	<i>Handroanthus chrysotrichus</i> (Mart. ex A. DC.) Mattos	N*	D			x		0.2
83	<i>Orbignya speciosa</i> (Mart. ex Spreng.) Barb. Rodr.	N*	PR	x				0.2
84	<i>Platymiscium pinnatum</i> (Jacq.) Dugand	N	D			x		0.2
85	<i>Tectona grandis</i> L. f.	E	D				x	0.2
86	<i>Alibertia edulis</i> (Rich.) A. Rich. ex DC.	N*	PR		x			0.2
87	<i>Guazuma tomentosa</i> Kunth	N*	PR		x			0.2
88	<i>Tapirira guianensis</i> Aubl.	N*	PR		x			0.2
89	<i>Vitex cymosa</i> Bertero ex Spreng.	N*	D		x			0.2
90	<i>Ficus gomelleira</i> Kunth & C.D. Bouché	E	D		x			0.2
91	<i>Machaerium aculeatum</i> Raddi	N*	D				x	0.2
92	Rubiaceae sp.	-	-				x	0.2
93	<i>Sterculia striata</i> A. St.-Hil. & Naudin	N*	D				x	0.2
94	<i>Ficus insipida</i> Willd.	N	D		x			0.1
95	<i>Pseudobombax longiflorum</i> (Mart.) A. Robyns	N*	D				x	0.1
96	<i>Spondias dulcis</i> Parkinson	E	D		x			0.1
97	<i>Syzygium jambolanum</i> (Lam.) DC.	E	PR		x			0.1
98	<i>Acosmium subelegans</i> (Mohlenbr.) Yakovlev	N*	D				x	0.1
99	<i>Annona crassiflora</i> Mart.	N*	D		x			0.1
100	<i>Aspidosperma subincanum</i> Mart. ex A. DC.	N*	D				x	0.1
101	<i>Bauhinia variegata</i> L.	N*	D			x		0.1
102	<i>Citrus</i> sp. 3	E	PR		x			0.1
103	<i>Hirtella gracilipes</i> (Hook. f.) Prance	N*	D		x			0.1
104	<i>Machaerium acutifolium</i> Vogel	N*	PR				x	0.1
105	<i>Pouteria gardneri</i> (Mart. & Miq.) Baehni	N*	D		x			0.1
106	<i>Zanthoxylum rhoifolium</i> Lam.	N*	PR		x			0.1
107	<i>Apeiba tibourbou</i> Aubl.	N*	PR				x	0.1
108	<i>Aspidosperma cylindrocarpon</i> Muell. Arg.	N*	PR				x	0.1
109	<i>Brosimum gaudichaudii</i> Trécul	N*	D		x			0.1
110	<i>Cordia ecalyculata</i> Vell.	N	D		x			0.1
111	<i>Cydistax antisiphilitica</i> (Mart.) Mart.	N*	D			x		0.1
112	<i>Eriotheca gracilipes</i> (K. Schum.) A. Robyns	N*	D				x	0.1
113	<i>Inga edulis</i> Mart.	N*	PR				x	0.1

Table 1. Continued...

N	Species, genus, family or absence of classification	Or.	Dec.	Groups				RAb.
				P	F	O	Oth.	
114	<i>Maclura tinctoria</i> (L.) D. Don ex Steud.	N*	D		x			0.1
115	<i>Morinda citrifolia</i> L.	E	PR				x	0.1
116	<i>Persea americana</i> Mill.	N	PR		x			0.1
117	<i>Vitex polygama</i> Cham.	N*	D		x			0.1
118	<i>Xilopia aromatica</i> Mart.	N*	D		x			0.1
119	<i>Caesalpinia echinata</i> Lam.	N	D				x	0.1
120	<i>Eucalyptus</i> sp. 4	E	D				x	0.1
121	<i>Plinia cauliflora</i> (DC.) Kausel	N	D		x			0.1
122	<i>Pseudobombax tomentosum</i> (Mart. & Zucc.) A. Robyns	N*	D				x	0.1
123	<i>Rhamnidium elaeocarpum</i> Reissek	N*	D		x			0.1
124	<i>Schizolobium parahyba</i> var. <i>amazonicum</i> (Huber ex Ducke) Barneby	N	D				x	0.1
125	<i>Trema micrantha</i> (L.) Blume	N*	D		x			0.1
126	<i>Adenantha pavonina</i> L.	E	PR				x	0.1
127	<i>Annona coriacea</i> Mart.	N*	PR		x			0.1
128	<i>Azadirachta indica</i> A. Juss.	E	PR				x	0.1
129	<i>Bauhinia</i> sp.	N*	D			x		0.1
130	<i>Caesalpinia ferrea</i> Mart. ex Tul.	N*	PR				x	0.1
131	<i>Caryocar villosum</i> (Aubl.) Pers.	N	D		x			0.1
132	<i>Copernicia australis</i> Becc.	N*	PR	x				0.1
133	<i>Eucalyptus</i> sp. 3	E	D				x	0.1
134	<i>Plathymenia reticulata</i> Benth.	N*	D				x	0.1
135	<i>Pouteria ramiflora</i> (Mart.) Radlk.	N*	D		x			0.1
136	<i>Spondias mombin</i> L.	N*	D		x			0.1
137	<i>Calophyllum brasiliense</i> Cambess.	N*	D		x			**
138	<i>Carica</i> sp.	E	PR		x			**
139	<i>Citrus</i> sp.	E	PR				x	**
140	<i>Dilodendron bipinnatum</i> Radlk.	N*	D		x			**
141	<i>Eugenia emarginata</i> (Kunth) DC.	N*	PR		x			**
142	Fabaceae sp.	-	-				x	**
143	<i>Guarea silvatica</i> C. DC.	N	PR				x	**
144	<i>Handroanthus ochraceus</i> (Cham.) Mattos	N*	D			x		**
145	<i>Lagenaria siceraria</i> (Molina) Standl.	E	D				x	**
146	<i>Magnolia grandiflora</i> L.	E	PR			x		**
147	Malvaceae sp.	-	-				x	**
148	<i>Moringa oleifera</i> Lam.	E	PR				x	**
149	<i>Ochroma pyramidale</i> (Cav. ex Lam.) Urb.	N	PR				x	**
150	<i>Platymiscium floribundum</i> Vogel	N*	PR				x	**
151	<i>Pouteria caimito</i> (Ruiz & Pav.) Radlk.	N*	PR		x			**
152	<i>Qualea multiflora</i> Mart.	N*	D				x	**
153	<i>Qualea parviflora</i> Mart.	N*	D				x	**
154	<i>Roystonea oleracea</i> (Jacq.) O.F. Cook	E	PR	x				**
155	<i>Sapium lanceolatum</i> Huber	N*	D		x			**
156	<i>Spathodea campanulata</i> P. Beauv.	E	PR			x		**

Table 1. Continued...

N	Species, genus, family or absence of classification	Or.	Dec.	Groups				RAb.
				P	F	O	Oth.	
157	<i>Syzygium cumini</i> (L.) Skeels	E	D		x			**
158	<i>Triplaris americana</i> L.	N*	PR			x		**
159	<i>Annona reticulata</i> L.	E	PR		x			**
160	<i>Attalea phalerata</i> Mart. ex Spreng.	N*	PR	x				**
161	<i>Averrhoa carambola</i> L.	E	PR		x			**
162	<i>Bauhinia cuiabensis</i> Steud.	E*	D			x		**
163	<i>Bauhinia rufa</i> (Bong.) Steud.	E*	D			x		**
164	<i>Bauhinia</i> sp. 2	E*	D			x		**
165	<i>Bauhinia</i> sp. 3	E*	D			x		**
166	<i>Bertholletia excelsa</i> Bonpl.	N	D		x			**
167	<i>Bixa orellana</i> L.	N*	PR				x	**
168	<i>Byrsonima coccolobifolia</i> Kunth	N*	D		x			**
169	<i>Capparis retusa</i> Griseb.	E	PR		x			**
170	<i>Citrus</i> sp. 2	E	PR				x	**
171	<i>Coccoloba mollis</i> Casar.	N*	D		x			**
172	<i>Copernicia prunifera</i> (Mill.) H.E. Moore	N*	PR	x				**
173	<i>Dracaena</i> sp.	E	PR				x	**
174	<i>Erythrina mulungu</i> Mart. ex Benth.	E	D			x		**
175	<i>Eugenia bimarginata</i> DC.	N*	PR		x			**
176	Euphorbiaceae sp.	-	-				x	**
177	<i>Garcinia gardneriana</i> (Planch. & Triana) Zappi	N*	PR		x			**
178	Indeterminada 1	-	-				x	**
179	Indeterminada 2	-	-				x	**
180	<i>Jatropha curcas</i> L.	E*	D				x	**
181	<i>Khaya ivorensis</i> A. Chev.	E	PR				x	**
182	Malvaceae sp. 2	-	-				x	**
183	<i>Pinus caribaea</i> Morelet	E	PR				x	**
184	<i>Plumeria rubra</i> L.	E	D			x		**
185	<i>Psidium cattleianum</i> Sabine	N*	PR		x			**
186	<i>Psidium</i> sp.	E	D		x			**
187	<i>Roupala montana</i> Aubl.	N*	D				x	**
188	<i>Schefflera morototoni</i> (Aubl.) Maguire, Steyerl. & Frodin	N*	PR		x			**
189	<i>Stryphnodendron adstringens</i> (Mart.) Coville	N*	D				x	**
190	<i>Zygia</i> sp.	N*	D				x	**
Total		-	-	7	71	32	80	100

Origin (Or.): Native to Brazil (N), Exotic (E). Natural occurrence in the phytogeographical domain of the Cerrado (*); Deciduity (Dec.): Deciduous (D), Perennial (PR); Groups: Palm trees (P), Fruitful (F), Ornamental (O), Other (Oth.); Value < 0.1 (**).

The values of the twelve most frequent species represent 53.2% of all individuals found on campus. Milano & Dalcin (2000) consider that the quantity of individuals of a species should not exceed 15% of the total population. In this case, the species from the UFMT fall within the recommendations of these authors.

Regarding deciduity, 64.7% of the species were deciduous, 31.6% perennial and 3.7% were not classified due to lack of identification. When compared to the proportion of autochthonous species on campus, this result can be explained since the majority of naturally occurring Cerrado species are characterized by leaf loss in the dry season. However,

Cuiabá is a city with high temperatures throughout the year, especially during August to October, so that it is necessary to implant perennial species such as *Licania tomentosa*, the most abundant species in the afforestation of the campus. For Lorenzi (2008), the preference for this species in the composition of roadside trees may be justified by its dense and perennial crown, which provides excellent shading and a deep root system. Therefore, its use is adequate to help mitigate the effect of the heat islands reported in the city of Cuiabá in other studies (Alves & Biudes, 2011; Santos et al., 2014).

Makhelouf (2009) states that the decreasing temperatures of the parks promote a reduction of the urban heat island, and points to green spaces as being less polluted places in a city. According to the author, green areas have a significant contribution to improving the microclimate and reducing the pollution rate in cities, and denominates these sites as “a city’s lungs”, being the most important planning sites in the city.

In an interview conducted by Batista et al. (2014), 39% of people are motivated to move to the UFMT because of the zoo, followed by a 33% group that seeks the vast green area. Also, campus-goers described a sense of well-being that favors physical and mental health, provided by the wide presence of trees. The environmental perception was also evaluated in the same study, which reported a feeling of silence, satisfaction, tranquility and peace found by the regulars.

Of the total species, 34.7% produce fleshy fruits, 16.6% show intense flowering, 3.7% are palm trees and 42.1% were in the species group of environmental services. Although they are allochthonous species to the region, *Mangifera indica*, *Genipa americana*, *Psidium guajava* and *Anacardium occidentale* are trees that produce attractive fleshy fruits to the avifauna of Cuiabá and mainly for the UFMT zoo, which naturally complements their diet and helps in the seed dispersal process. *Dipteryx alata*, *Samanea tubulosa*, *Guazuma ulmifolia*, *Caryocar brasiliense*, *Buchenavia tomentosa*, *Casearia sylvestris* and others are also indigenous species found on the UFMT campus, which contribute to maintaining diversity.

Pivetta & Silva Filho (2002) indicate that one of the advantages that trees show by their natural characteristics in the urban environment is to provide a better aesthetic effect. The *Handroanthus* genus comprises about 9.1% of all campus individuals, and is widely recommended and used in urban tree growing due to the beauty of its flowering. *Bauhinia* is another genus found on campus with a few specimens; however, it is much used in urban afforestation both for the beauty of its flowers and for its small to medium size, which allows its use in roads under the electricity grid. Palm trees are widely used in the UFMT, where almost all their abundance is concentrated in the central road beds. However, its use is

considered inadequate in road forestation and its planting in the urban zone of Cuiabá is prohibited, which includes the UFMT campus (Cuiabá, 2012). Thus, its substitution by woody species is indicated.

At the UFMT, 42.1% of the species were classified as palm trees, fruit trees or ornamental, which were classified as environmental service providers. Morgenroth et al. (2016) argue that urban trees are important because they regulate the environment in many ways in the city, such as reducing air pollution and storm water management. For Clapp et al. (2014), these environmental services provided by trees in the urban environment depend on their specific characteristics, namely morphological, physiological, temporal, and seasonal or diurnal effects of different species. In this sense, the urban afforestation as a whole must be planned with different species, since each one has its particularities and should be used in the search for environmental balance.

Of the total number of species present on campus, 72.6% are native to Brazil, 23.7% originate in other countries and 3.7% are undetermined due to the non-botanical identification of the species. For the species native to Brazil, *Licania tomentosa* and *Jacaranda cuspidifolia* stand out for having the largest number of individuals. Among the species exotic to Brazil, *Mangifera indica* and *Psidium guajava* stood out with greater abundance. In addition to the majority of native species to Brazil, 62.6% of these species occur naturally in the phytogeographic domains of the Cerrado. Despite having an approximate value as required, UFMT afforestation must comprise at least 70% native species (Cuiabá, 2012). For Farias Castro et al. (2011), the use of autochthonous species in urban afforestation has great potential for both its adaptation to the environment and for preserving biodiversity; both of which are fundamental for balancing the local environment.

According to Riley et al. (2018), the contribution of exotic species to providing ecosystem services should be considered when in appropriate proportions, because, just as with native species, they also perform a number of important benefits in cities, such as carbon sequestration, heat island mitigation, air quality improvement, surface runoff reduction, increased property values, improvement of human health and well-being, and promotion of a greater sense of social connectivity. For Morgenroth et al. (2016), allochthonous species are more likely to positively respond to fluctuating environmental conditions than native species, so it is advantageous to use them.

Based on data from the Instituto Hórus (Instituto Hórus de Desenvolvimento e Conservação Ambiental, 2017), it was found that some exotic species used in the UFMT campus afforestation are characterized by rapid propagation and invasive potential, such as *Albizia lebbbeck*, *Leucaena*

leucocephala, *Artocarpus heterophyllus*, *Azadirachta indica*, *Clitoria fairchildiana*, and others. For Ziller (2001), invasive species can have disastrous impacts on the environment and directly affect agricultural, forestry or ornamental activities. In this context, it is necessary to monitor the propagation of these specimens given their rapid development, which may promote greater competition with native species and their possible suppression.

The diversity obtained by the Shannon Index (H') was 4.1, being considered high since it varies from 0 to 5, in which the higher the value, the greater the diversity. The index is based on the number of species present in a given environment, added to the uniformity between them and their relation to the abundance and dominance degree of each one (Odum, 2012). The calculated value for the Simpson Diversity Index was 0.04, which expresses high diversity and corroborates the result of the estimated Shannon-Weaver Index. This index expresses the probability that two randomly selected individuals belong to the same species and their value varies from 0 to 1; the higher the value, the greater the dominance of certain species; on the other hand, the lower the diversity.

Biondi & Lima Neto (2011) point out the importance of vegetal species diversification in the urban environment as a source for establishing richer and more diversified fauna. Morgenroth et al. (2016) reiterate the need for species diversity to achieve a stock of resilient trees to contribute to the urban ecosystem stability, where all tree species have desirable and undesirable characteristics, but species selection must be performed strategically to optimize the ecosystem services. Santos et al. (2014) indicate that the green areas in Cuiabá have specific microclimate characteristics, which are different from the non-wooded areas, with a higher formation of humidity islands; a fact that the authors attributed to the consequent decrease in temperature and an increased relative humidity of these places. A similar result was obtained by Alves & Biudes (2011) in a study on the microclimate on the UFMT campus. However, Marandola Junior et al. (2011) warn about reducing areas for permanent preservation of the streams of Cuiabá; a fact that contributes to an unstable and fragile urban ecosystem and potentiates the risks arising from floods in the capital of Mato Grosso. In this context, the campus green area's contribution to the quality of life of the population regarding mitigation of high temperatures, lowering the low relative humidity and increasing the water flow of the soil of Cuiabá, is corroborated.

The green area of the UFMT is very important for maintaining fauna and flora diversity in the municipality of Cuiabá and can be a reference for planning and implementing other green areas. Many of the local fauna species depend on the food

production provided by the fruit species in the campus, as well as the seed dispersion of these vegetal species depending on the fauna displacement for propagating its descendants.

4. CONCLUSIONS

The afforestation of the UFMT campus in Cuiabá is rich, with great species diversity in an adequate proportion of Brazilian native species. However, the floristic composition needs adjustment to the native species in the phytogeographic domain of the Cerrado. The floristic structure occasionally needs to be adjusted regarding exotic species with invasive potential, replacing them with autochthonous species. The implanted species in the roadside tree planting must be adapted to the municipal afforestation legislation.

ACKNOWLEDGEMENTS


To the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes) for granting a Graduate scholarship to the first author during the study. To UFMT along with PET/Floresta for the opportunity to research and contribute to the knowledge on campus afforestation. To Professor Dr. Diego Tyszka Martinez for the enriching guidance and contribution to the study.

SUBMISSION STATUS

Received: 29 Mar. 2018

Accepted: 13 Sept. 2018

Associate editor: Rodrigo Studart Corrêa

 0000-0002-9422-2629

CORRESPONDENCE TO

Poliane Pierra Patricio Pazini

Universidade Federal de Mato Grosso (UFMT), Av. Fernando Correa da Costa, 2367, CEP 78060-900, Cuiabá, MT, Brasil

e-mail: polianepatricio@gmail.com

FINANCIAL SUPPORT

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes), Grant/Award number: 36000,00.

REFERENCES

- Alencar LD, Cardoso JC. Paisagismo funcional: o uso de projetos que integram mais que ornamentação. *Ciência, Tecnologia e Ambiente* 2015; 1(1): 1-7.
- Alvares CA, Stape JL, Sentelhas PC, Moraes Gonçalves JL, Sparovek G. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift* 2013; 22(6): 711-728. 10.1127/0941-2948/2013/0507
- Alves EDL, Biudes MS. Os microclimas da Universidade Federal de Mato Grosso/Cuiabá. *Raega* 2011; 23: 600-620. 10.5380/raega.v23i0.24923

- Batista BMF, Sousa RAT, Zavala AAZ, Pasa MC, Timofeiczuk Junior R. Percepção ambiental dos frequentadores do campus da Universidade Federal de Mato Grosso. *Flovet* 2014; 1(6): 98-114.
- Biondi D. Floresta urbana: conceitos e terminologias. In: Biondi D. *Florestas urbanas*. Curitiba: Produção Independente; 2015. p. 11-28.
- Biondi D, Lima Neto EM. Introdução de espécies na arborização de ruas. In: Biondi D, Lima Neto EM. Pesquisa em arborização de ruas. Curitiba: Produção Independente, 2011. p. 9-28.
- Clapp JC, Ryan III HDP, Harper RW, Bloniarz DV. Rationale for the increased use of conifers as functional green infrastructure: a literature review and synthesis. *Arboricultural Journal* 2014; 36(3): 161-178. 10.1080/03071375.2014.950861
- Cuiabá. Decreto n. 5.144, de 15 de fevereiro de 2012. Dispõe sobre a arborização pública na área urbana e sedes de distrito do município de Cuiabá e dá outras providências. *Palácio Alencastro*, Cuiabá (2012 Feb. 15).
- Eisenlohr PV, Carvalho-Okano RM, Vieira MF, Leone FR, Stringheta ACO. Flora fanerogâmica do campus da Universidade Federal de Viçosa, Viçosa, Minas Gerais. *Ceres* 2008; 55(4): 317-326.
- Farias Castro AS, Moro MF, Rocha FCL. Plantas dos espaços livres da reitoria da Universidade de Fortaleza (Unifor), Ceará, Brasil. *Revista Brasileira de Biociências* 2011; 9(1): 126-129.
- Fundação Coordenação de Projetos, Pesquisas e Estudos Tecnológicos – Coppetec. *Flora do Brasil 2020*. Rio de Janeiro; c2019 [cited 2017 Jan 25]. Available from: <http://bit.ly/2NoEVy0>
- Instituto Brasileiro de Geografia e Estatística – IBGE. *Estimativas da população residente no Brasil e unidades da federação com data de referência em 1º de julho de 2017*. Brasília, DF; 2017.
- Instituto Hórus de Desenvolvimento e Conservação Ambiental. *Base de dados nacional de espécies exóticas invasoras I3N Brasil*. Florianópolis; 2017 [cited 2017 Jan 25]. Available from: <http://bit.ly/34g76WG>
- Kramer JA, Krupek RA. Caracterização florística e ecológica da arborização de praças públicas do município de Guarapuava, PR. *Árvore* 2012; 36(4): 647-658. 10.1590/S0100-67622012000400007
- Loboda CR, Angelis BLD. Áreas verdes públicas urbanas: conceitos, usos e funções. *Ambiência* 2005; 1(1): 125-139.
- Lorenzi H. *Árvores brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil*. Nova Odessa: Plantarum, 2008. v. 3.
- Makhelouf A. The effect of green spaces on urban climate and pollution. *Journal of Environmental Health Science and Engineering* 2009; 6(1): 35-40.
- Marandola Junior E, D'Antona AO, Ojima R. *População, ambiente e desenvolvimento: mudanças climáticas e urbanização no centro-oeste*. Campinas: Núcleo de Estudos da População/Unicamp; 2011.
- McHale MR, McPherson EG, Burke IC. The potential of urban tree plantings to be cost effective in carbon credit markets. *Urban Forestry and Urban Greening* 2007; 6: 49-60. 10.1016/j.ufug.2007.01.001
- Milano MS, Dalcin EC. *Arborização de vias públicas*. Rio de Janeiro: Light; 2000.
- Missouri Botanical Garden. *Tropicos*. Saint Louis; 2016 [cited 2017 Jan 1]. Available from: <http://bit.ly/36f9TkG>
- Morgenroth J, Östberg J, van den Bosh CK, Nielsen AB, Hauer R, Sjöman H et al. Urban tree diversity: taking stock and looking ahead. *Urban Forestry and Urban Greening* 2016; 15: 1-5. 10.1016/j.ufug.2015.11.003
- Odum EP. *Ecologia*. Rio de Janeiro: Guanabara Koogan; 2012.
- Pinheiro CR, Souza DD. A importância da arborização nas cidades e sua influência no microclima. *Gestão e Sustentabilidade Ambiental* 2017; 6(1): 67-82. 10.19177/rgsa.v6e1201767-82
- Pivetta KFL, Silva Filho DF. *Arborização urbana*. Jaboticabal: Unesp/FCAV/Funep; 2002. (Boletim Acadêmico: Série Arborização Urbana). 74 p.
- R Development Core Team. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2015.
- Riley CB, Herms DA, Gardiner MM. Exotic trees contribute to urban forest diversity and ecosystem services in inner-city Cleveland, OH. *Urban Forestry and Urban Greening* 2018; 29: 367-376. 10.1016/j.ufug.2017.01.004
- Santos HG, Jacomine PKT, Anjos LHC, Oliveira VA, Lumbreiras JE, Coelho MR et al. *Sistema brasileiro de classificação de solos*. 3rd ed. Brasília, DF: Embrapa Solos, 2013.
- Santos FMM, Oliveira AS, Nogueira MCJA, Musis CJ, Nogueira JS. Análise do clima urbano de Cuiabá, MT, Brasil por meio de transectos móveis. *Paranoá* 2014; (11): 45-53. 10.18830/issn.1679-0944.n11.2014.12083
- Stevens PF. Angiosperm Phylogeny Website, 2016 [cited 2016 Sept. 10]. Available from: <http://bit.ly/2N4oU1e>
- Ziller SR. Plantas exóticas invasoras: a ameaça da contaminação biológica. *Ciência Hoje* 2001; (178): 77-79.