

Current status of the tomato pinworm, *Tuta absoluta* (Meyrick) in the Indian sub-continent: Challenges ahead

K.S. Nitin and A. K. Chakravarthy

Society for Science and Technology Applications (SSTA)
(Formerly, Head, Divn. of Entomology and Nematology, IIHR)
No. 7, 3rd cross, Chowdaiah block, R.T.Nagar, Bangalore -560032 India.
Corresponding author: chakravarthyakshay@gmail.com

The tomato pinworm, *Tuta absoluta* is an oligophagous pest, infesting economically important solanaceous crops (Sridhar *et al.*, 2014; Nitin *et al.*, 2019). In India, for the first time it was reported on tomato from the south Indian state, Karnataka (Sridhar *et al.*, 2014). Rapidly it invaded all the other south Indian states *viz.* Telangana (Kumari *et al.*, 2015), Tamil Nadu (Shanmugam *et al.*, 2016), Kerala (Sivakumar *et al.*, 2017), Andhra Pradesh (Rasheed *et al.* 2018) and North Indian states *viz.*, Maharashtra (Shashank *et al.*, 2015), Gujarat (Chavan *et al.*, 2016), New Delhi (Shashank *et al.*, 2016), Madhya Pradesh (Swathi *et al.*, 2017), Punjab (Sandeep *et al.*, 2017), Meghalaya (Sankarganesh *et al.*, 2017), Himachal Pradesh (Sharma and Gavkare 2017), Uttarakhand (Singh and Panchbhaiya 2018), Chhattisgarh (Balaji *et al.*, 2018), Uttar Pradesh (Halder *et al.*, 2019), Bihar and Odisha (Sridhar and Srinivas 2019) (Fig. 1). This indicates that *T. absoluta* has now established itself in almost all parts of India in just about 5 years. This could be attributed to the un-regulated transportation of solanaceous fruits and plants especially tomatoes across the country. There are no reports of *T. absoluta* infestations from Rajasthan, Jammu and Kashmir. This may be because of upheaval climato-geographic conditions of these regions and solanaceous plants not being major crops there. In Jharkhand and other north eastern states extensive surveys could perhaps confirm *T. absoluta* infestation as these regions possess favourable conditions during certain months of the year.

The tomato pinworm, *T. absoluta* has the potential to occur throughout the year in Indian agro-climatic conditions (Nitin *et al.*, 2017; Nayana *et al.*, 2018). Their activity and population increase with the increase in atmospheric temperature upto certain extent, (36°C) (Nitin *et al.*, 2018), with significant damage to tomato production. Researchers have investigated COI gene sequences of *T. absoluta* and found that it is 100% homologous with the sequences deposited by researchers in NCBI database (Sankarganesh *et al.*, 2017; Shashank *et al.*, 2018), indicating

spread of *T. absoluta* from a single source. Genetic variations in *Tuta* populations are not yet detected.

Indian farmers are using pheromone traps for the early detection, monitoring and mass trapping of *T. absoluta* in their fields (Bhanu *et al.*, 2017). Along with the pheromone traps, solar traps also give promising results by trapping female *T. absoluta* (Sridhar *et al.*, 2019). Farmers are using synthetic insecticides as a primary control measure for *T. absoluta* management in spite of its drawbacks like environmental pollution, non-target effects and chance of developing insecticide resistance by the pest. Chemicals such as spinetoram, cyantraniliprole, flubendiamide and spinosad have given promising results in managing this pest both during vegetative and fruiting stages (Sridhar *et al.*, 2016). In some countries *T. absoluta* has developed resistance to diamide (Roditakis *et al.*, 2015), chlorantraniliprole (Silva *et al.*, 2016) and organophosphate and pyrethroid insecticides (Zibae *et al.*, 2018), but in India *T. absoluta* showed reduced susceptibility to certain commonly used insecticide (Prasanna kumar *et al.*, 2020). *Tuta absoluta* management failure by chemical control is predicted to appear in future leading to adoption of an integrated pest management approach comprising eco-friendly methods. Integrated pest management approaches comprising of pheromones, microbials, natural enemies and eco-friendly molecules are more promising in controlling *T. absoluta* both environmentally and economically (Ballal *et al.*, 2016; Nitin *et al.*, 2018; Buragohain *et al.*, 2020). Along with these trials, researcher also found that the usage of yellow incandescent bulb traps to be very effective in attracting *T. absoluta* followed by blue light traps that captures both males and females (Sridhar *et al.*, 2019).

For the successful management of this noxious pest, Indian researchers should concentrate towards developing commercial genetically modified (GM) crop varieties that are resistant to *T. absoluta*, exploiting novel techniques such as RNA interference (RNAi) to silence the genes that result in impaired growth and/or development and even mortality of the pest. Sterile insect technique (SIT) can be employed for the management in a mass region as it is species-specific, environmentally friendly, and can be combined with other management options. Along with these interventions, legislations also play an important role in the management of invasive pests. Listing *T. absoluta* under quarantine species list, implementing cross-border quarantine facilities across the states, bringing public awareness through campaigns, conducting

farmers meets with the experts and sharing the knowledge among the stakeholders help in managing this invasive pest. Frequent exposure to insecticides, short lifecycle and generation period and higher rates of invasiveness are challenges to come in future.

References

- Balaji, D. R., Jeyarani, S., Ramaraju, K., Mohankumar, S. and Shanmugam, P. S. 2018. Occurrence of South American tomato pinworm, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae): An invasive pest in Tamil Nadu, India. *Journal of Entomology and Zoology Studies*, 6(2), 657-662.
- Ballal, C. R., Gupta, A., Mohan, M., Lalitha, Y. and Verghese, A. 2016. The new invasive pest *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in India and its natural enemies along with evaluation of Trichogrammatids for its biological control. *Current Science*, 2155-2159.
- Bhanu, K. R. M., Ramachandra, V. A. W. and Mamatha, B. 2017. Monitoring of *Tuta absoluta* (tomato leaf miner) using pheromone traps: a recent invasive pest in India. *Environment and Ecology*, 35(4A), 2971-2976.
- Buragohain, P., Saikia, D. K., Sotelo-Cardona, P. and Srinivasan, R. 2020. Development and validation of an integrated pest management strategy against the invasive South American tomato leaf miner, *Tuta absoluta* in South India. *Crop Protection*, 139, 105348.
- Chavan, S.M., Pravinkumar, M., and Pandya, C. D. 2016. South American Tomato Pinworm, *Tuta absoluta*: A New Invasive Insect Pest Recorded on Tomato in Gujarat. *Advances in Life Sciences* 5(15), 5736-5738.
- Halder, J., Kushwaha, D., Rai, A. B. and Singh, B. 2019. Biology and Biorational Management of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae): A Global Challenge to Tomato Production. In *Proceedings of the Zoological Society* (Vol. 72, No. 2, pp. 107-110). Springer India.

- Kumari, D. A., Anitha, G., Anitha, V., Lakshmi, B. K. M., Vennila, S. and Rao, N. H. P. 2015. New record of leaf miner, *Tuta absoluta* (Meyrick) in Tomato. *Insect Environment*, 20(4), 136-138.
- Nayana, B. P., Shashank, P. R. and Kalleshwaraswamy, C. M. 2018. Seasonal incidence of invasive tomato leaf miner, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) on tomato in Karnataka, India. *Journal of Entomology and Zoology Studies*, 6(1), 400-405.
- Nitin, K. S., Chakravarthy, A. K., Özgökçe, M. S. and Atlihan, R. 2019. Population growth potential of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) on tomato, potato, and eggplant. *Journal of Applied Entomology*, 143(5), 518-526.
- Nitin, K. S., Naik, O. S., Sridhar, V., Bhat, P. S. and Chakravarthy, A. K. 2018. Development and validation of Integrated Pest Management modules against South American tomato moth, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in India: Blending non-insecticidal, nature friendly tools. *Pest Management in Horticultural Ecosystems*, 24(2), 115-120.
- Nitin, K. S., Sridhar, V., Kumar, K. P. and Chakravarthy, A. K. 2017. Seasonal incidence of South American tomato moth, *Tuta absoluta* (Meyrick)(Gelechiidae: Lepidoptera) on tomato ecosystem. *Int. J. Pure. Appl. Bioscie*, 5, 521-525.
- Nitin, K. S., Vaddi, S., Naik, S. O., Chakravarthy, A. K. and Atlihan, R. 2018. Effect of temperature and CO₂ on population growth of South American Tomato Moth, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) on tomato. *Journal of Economic Entomology*, 111(4), 1614-1624.
- Prasannakumar, N. R., Jyothi, N., Saroja, S. and Kumar, G. R. 2020. Relative toxicity and insecticide resistance of different field population of tomato leaf miner, *Tuta absoluta* (Meyrick). *International Journal of Tropical Insect Science*, 1-9.

- Rasheed, V. A., Rao, S. K., Babu, T. R., Murali, T., Krishna, B. V. and Naidu, G. M. 2018. Incidence of South American tomato leaf miner, *Tuta absoluta* (Meyrick) in Chittoor district of Andhra Pradesh, India.
- Roditakis, E., Vasakis, E., Grispou, M., Stavrakaki, M., Nauen, R., Gravouil, M. and Bassi, A. 2015. First report of *Tuta absoluta* resistance to diamide insecticides. *Journal of Pest Science*, 88(1), 9-16.
- Sandeep K.S, Sridhar V, Sharma A. and Asokan R. 2017. Report on the occurrence of South American Tomato moth, *Tuta absoluta* (Meyrick) in Punjab, India as evident from trap catches and molecular diagnosis. *Pest Manag. Hortil. Ecosyst.* 23(1):89-91.
- Sankarganesh, E., Firake D.M., Sharma, B, Verma, V.K and Behere G.T. 2017. Invasion of the South American Tomato Pinworm, *Tuta absoluta*, in northeastern India: a new challenge and biosecurity concerns. *Entomol. Gen* 2017;36(4):335-345.
- Shanmugam, P. S., Ramaraju, K. and Indhumathi, K. 2016. First record of South American tomato moth, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae) in Tamil Nadu, India. *Entomon*, 41(1), 61-66.
- Sharma PL and Gavkare O. 2017. New distributional record of invasive pest *Tuta absoluta* (Meyrick) in North-Western Himalayan Region of India. *Nat. Acad. Sci. Lett. Springer*.
- Shashank P.R., Suroshe S.S., Singh P.K., Chandrashekar K, Nebapure S.M. and Meshram N.M. 2016. Report of invasive tomato leaf miner, *Tuta absoluta* (Lepidoptera: Gelechiidae) from northern India. *Indian J Agric. Sci.* 86(12):1635-1636.
- Shashank, P. R., Chandrashekar, K., Meshram, N. M. and Sreedevi, K. 2015. Occurrence of *Tuta absoluta* (Lepidoptera: Gelechiidae) an invasive pest from India. *Indian Journal of Entomology*, 77(4), 323-329.
- Shashank, P. R., Twinkle, S., Chandrashekar, K., Meshram, N. M., Suroshe, S. S. and Bajracharya, A. S. R. 2018. Genetic homogeneity in South American tomato pinworm, *Tuta absoluta*: a new invasive pest to oriental region. *3 Biotech*, 8(8), 350.

- Silva, J. E., Assis, C. P., Ribeiro, L. M. and Siqueira, H. A. 2016. Field-evolved resistance and cross-resistance of Brazilian *Tuta absoluta* (Lepidoptera: Gelechiidae) populations to diamide insecticides. *Journal of economic entomology*, 109(5), 2190-2195.
- Singh, D. K. and Panchbhैया, A. 2018. First record of tomato leaf miner, an invasive pest in Uttarakhand, India under polyhouse condition. *Journal of Hill Agriculture*, 9(1), 127-130.
- Sivakumar, T., Josephraj Kumar, A. and Anitha, N. 2017. First report of tomato pinworm, *Tuta absoluta* (Meyrick) on eggplant *Solanum melongena* L. from Kerala, India. *Entomon*, 42(4), 335-338.
- Sridhar V, Chakravarthy AK, Asokan R, Vinesh LS, Rebijith KB, Vennila S. 2014. New record of the invasive South American tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in India. *Pest Manag Hortic Ecosyst*.20:148-154.
- Sridhar, V. and Srinivas, P. 2019) Report of South American tomato moth, *Tuta absoluta* (Meyrick) from Odisha. *Pest Management in Horticultural Ecosystems*, 25(1), 119-120.
- Sridhar, V., Onkaranaik, S. and Nitin, K. S. 2016. Efficacy of new molecules of insecticides against South American tomato moth, *Tuta absoluta* (Meyrick)(Lepidoptera: Gelechiidae). *Pest Management in Horticultural Ecosystems*, 22(2), 137-145.
- Sridhar, V., Naik, S. O., Nitin, K., Asokan, R., Swathi, P. and Gadad, H. 2019. Efficacy of integrated pest management tools evaluated against *Tuta absoluta* (Meyrick) on tomato in India. *Journal of Biological Control*, 33(3), 264-270.
- Swathi, P., Swathi, B., Das, S. B., Sridhar, V., Giribabu, O., Snehalatha, G. and Raypuriya, N. 2017. First report of South American tomato leaf miner, *Tuta absoluta* (Meyrick) from Madhya Pradesh, India. *Pest Management in Horticultural Ecosystems*. 23(1):92-93.
- Zibae, I., Mahmood, K., Esmaily, M., Bandani, A. R. and Kristensen, M. 2018. Organophosphate and pyrethroid resistances in the tomato leaf miner *Tuta absoluta* (Lepidoptera: Gelechiidae) from Iran. *Journal of Applied Entomology*, 142(1-2), 181-191.



Fig. 4. *Tuta absoluta* infested tomato plant



Fig. 5. *Tuta absoluta* infested tomato fruit



Fig. 6. *Tuta absoluta* larval instars