

## **Genetic Characterization of Fall Armyworm (*Spodoptera frugiperda*, J.E. Smith) Feeding on Pearl Millet (*Pennisetum glaucum* L.R.Br.) in Tamil Nadu**

Sathyan T<sup>1</sup>, Sathiah N<sup>1\*</sup>, Mohankumar S<sup>2</sup>, Balasubramani V<sup>2</sup>, Kokiladevi E<sup>2</sup>,  
Ravikesavan R<sup>3</sup>, Srinivasan T<sup>3</sup> and Kennedy J S<sup>4</sup>

<sup>1</sup> Department of Agricultural Entomology, Centre for Plant Protection Studies, Tamil Nadu Agricultural University, Coimbatore – 641003, Tamil Nadu, India

<sup>2</sup> Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology Tamil Nadu Agricultural University, Coimbatore – 641003, Tamil Nadu, India

<sup>3</sup> Department of Millets, Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore – 641003, Tamil Nadu, India

<sup>4</sup> School of Post Graduate Studies, Tamil Nadu Agricultural University, Coimbatore – 641003, Tamil Nadu, India

\*Corresponding author's email id: nsathiah@gmail.com

### **Abstract**

In the present study, molecular characterization was carried out at the Department of Plant Biotechnology, Tamil Nadu Agricultural University, Coimbatore to identify the strains of fall armyworm collected from pearl millet crop at four different locations in Tamil Nadu, India. PCR-RFLP profile of mitochondrial *COI* fragment exhibited the presence of both 'C' and 'R' strains of FAW. Mt*COI* region analysis revealed that, thirteen samples showed 'R' strain identity, and others assumed 'C' strain identity. Sequence analyses of mt*COI* region of FAW feeding on pearl millet showed nucleotide variations in 22 positions. The Strain, ON142640 (Pearl millet Vridhachalam India) totally diverged from all populations subjected to phylogenetic analysis in this study and was suspected to be an inter-strain (R/C) hybrid. The

per cent identity analysis of *S. frugiperda* ranged from 99% to 100% with previously deposited sequences in the NCBI-GenBank database.

**Keywords:** *Fall armyworm; Pearl millet; Molecular characterization; Strain; PCR-RFLP; Phylogenetic analysis*

## Introduction

Pearl millet (*Pennisetum glaucum* L.R.Br.), popularly known as “Candle millet” or “Indian millet” grown widely in dry areas of the arid and semi-arid tropics where no other cereal can thrive (Reddy *et al.*, 2021). It covers 6.93 million hectares and produces an average of 8.61 million tonnes per year, with productivity of 1,243 kg ha<sup>-1</sup> (Directorate of Millets Development, 2020). Villupuram, Thoothukudi, Tiruvannamalai, Virudhunagar, Cuddalore, Vellore, and Madurai are the major pearl millet growing districts of Tamil Nadu (Season and Crop Report, 2022).

About 300 insects have been found feeding on pearl millet in various regions of the world (Sharma *et al.*, 1981), although the number that can cause major damage is likely less than a dozen, and those that cause serious damage on a continuous annual basis are even fewer (Verma, 1980). Recently, the invasive fall armyworm (FAW), *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) has been found infesting the pearl millet crop. Apart from maize, sorghum (60.1 per cent) was the most popular millet host, followed by pearl millet (41.4 per cent), barnyard millet (22.9 per cent), and finger millet (10.2 per cent) (ICAR-IIMR, 2019). FAW was reported on maize, sorghum, pearl millet, and finger millet in different districts of Andhra Pradesh from August to November, 2018 (Bhavani *et al.*, 2019). This pest's voracious feeding and long-distance flight behavior pose a serious threat to grain production. The pest feeds on more than 350 different plant species (Montezano *et al.*, 2018). Though it infests a wide range of plants, it has a clear affinity for graminaceous plants (Casmuz *et al.*, 2010).

The races in FAW can only be distinguished molecularly, not morphologically (Prowell *et al.*, 2004; Nagoshi *et al.*, 2007). FAW is genetically divided into two races: one is the corn-designated 'C' strain, which feeds mostly on maize, sorghum, cotton, and pulses, and the other is the rice-designated 'R' strain, which favors rice, sugarcane, and other short grasses (Pashley, 1998; Nagoshi *et al.*, 2016). Mitochondrial *COI* markers are frequently employed in insects to identify and determine population structure, and they are known for maternal inheritance, high copy number, conservative nature *etc.* (Mauro *et al.*, 2006). Identifying fall armyworm strains is aided by PCR-RFLP investigation of the mitochondrial

*COI* gene using restriction enzymes (Meager and Nagoshi , 2003; Nagoshi *et al.*, 2018). In this context, the current work used genetic approaches to identify the fall armyworm strain(s) feeding on pearl millet.

## **Materials and methods**

### **Insect collection and preservation**

FAW larvae (5 nos. in each location) were collected from pearl millet crop in four different locations and are listed in table 1. The collected insects were preserved individually in microfuge tubes and stored at -20 °C until taken for DNA extraction.

### **DNA preparation and quantification**

The genomic DNA of FAW was isolated using the CTAB technique (cetyl trimethylammonium bromide) (Doyle and Doyle, 1990). DNA was extracted from the prolegs of larvae. Tissues were homogenized thoroughly in 200 mL of extraction buffer (100 mM Tris-HCl, pH 8.0, 1.4M NaCl, 0.02M EDTA, two per cent CTAB, and 0.2 per cent mercaptoethanol), maintained at 65°C for an hour, then centrifuged for 15 minutes at 12,000 rpm. An equal volume of chloroform: isoamyl alcohol (24:1) was added to the supernatant and centrifuged at 10,000 rpm for 10 minutes to separate the phases. This process was repeated with the top aqueous phase transferred to a 1.5 µL sterile microfuge tube. About 200 µL of ice-cold isopropanol was added to this aqueous phase to precipitate the DNA, which was then incubated at -20 °C overnight. After incubation, the DNA pellet was obtained by centrifugation at 10,000 rpm for 10 minutes at 4 °C. The DNA pellet was washed twice with absolute ethanol and air-dried. Pellet was suspended in 30-50 µL of 1X TE buffer and stored at -20 °C until used for PCR analysis. The isolated DNA was analyzed quantitatively and qualitatively by Nanodrop Spectrophotometer (Genova, USA) and agarose gel electrophoresis, respectively.

### **PCR analysis**

The PCR amplification of DNA obtained from FAW larvae in various places was done. To run PCR, DNA samples were diluted with 1X TE buffer to obtain a working solution of 50-100 ng/L (working aliquot). The primers were synthesized from Bioserve Biotechnologies Private Limited, Hyderabad, India. The primer pair JM 76F (5'GAGCTGAATTAGG(G/A)ACTCCAGG3') and JM 77R (5'ATCACCTCC(A/T) CCTGCAGGATC3') were used to amplify the Mt*COI* region, yielding a 568-bp fragment. PCR amplification for all segments was carried out with 25 µL reaction mixture (12.5 µL of 2X

Thermo Scientific PCR Master Mix, 1 µL forward primer, 1 µL reverse primer, 9.5 µL nuclease-free water and 1 µL DNA template) using Eppendorf thermocycler with ambient thermo-cycling profile at 94 °C (1 min) followed by 33 cycles of 92 °C (45 s), 56 °C (45 s), 72 °C (1 min) and a final segment of 72 °C for 3 min. PCR amplified products were fractionated on a 2% PCR grade agarose horizontal gel and documented in the Bio-Rad Gel Doc XR imaging system.

### **RFLP analysis**

To determine strain identity, PCR products of the *COI* region were digested with *SacI* and *MspI* restriction enzymes purchased from Thermo Fisher Scientific and New England Biolabs, respectively. A reaction mixture including 10 µL PCR product, 2 µL buffer, 1 µL enzyme, and 17 µL nuclease-free water was prepared, incubated at 37 °C for 2 hours, fractionated on a 2% agarose gel, and documented.

### **Sequencing and GenBank submissions**

For double-pass DNA sequencing, the unpurified PCR amplified product (20 µL) was sent to Bioserve Biotechnologies India Pvt Ltd. in Hyderabad, Telangana, India. Bioedit software (version 7.25) was used to align and edit the raw sequence and construct the sequence identity and nucleotide difference count matrix. A dendrogram was created by aligning the select partial nucleotide sequences of *MtCOI* of *S. frugiperda* using Clustal W, and a phylogenetic tree was constructed using MEGA11 (version 11.0). The evolutionary distance between the sequences was determined using a bootstrap analysis with 1000 replications using the neighbor-joining method (Tamura *et al.*, 2021).

## **Results and discussion**

### **Strain identity analysis using PCR-RFLP**

The PCR analysis results revealed that the expected fragment size of about ~568 bp from *mtCOI* region was amplified in all the tested samples (Fig. 1A). Restriction enzymatic analysis of *mtCOI* segments exposed that, *SacI* enzyme cut the DNA into ~413 bp and ~155 bp fragments in Rice (R) strain whereas DNA remains uncut (~568 bp) in Corn (C) strain (Fig. 1B). Oppositely, *MspI* enzyme cut the DNA into ~461 bp and ~107 bp fragments in 'C' strain while DNA remains uncut (~568 bp) in 'R' strain (Fig. 1C). Out of 20, 13 samples showed 'R' strain identity, and seven assumed 'C' strain identity. The samples from Coimbatore and Vridhachalam comprised three corn and two rice strain individuals. Samples from Ramanathapuram possessed four rice and one corn strain. All the five samples collected from

Thalavaipuram assumed rice strain identity. Studies by Nagoshi *et al.* (2007) and Nagoshi and Meager (2004) implied that, the rice strain populations of FAW prefer to feed on millets, grass species *etc.* and corn strain populations prefer corn and sorghum for feeding.

Opposing results to findings were also known to occur. Irrespective of host plants, FAW reported on maize, sweet corn and sorghum assumed 'R' strain identity with minimal genetic diversity exhibiting no host/ location-specific variations (Swamy *et al.*, 2018). The presence of 'R' strain in all the FAW populations obtained from maize and sorghum crops in Swaziland was confirmed (Assefa, 2019). The FAW samples collected from Banten, Indonesia were assigned to the 'R' strain (Sartiami *et al.*, 2019).

### **Nucleotide variations**

Nucleotide variations were found at different places in the mtCOI region of FAW populations feeding on pearl millet (Fig. 2). The highest variation in the nucleotides was observed in OM462844 followed by OM465797 and OM463624 collected from pearl millet field at TNAU, Coimbatore, with 22, 21, and 5 bases, respectively. The strain from Vridhachalam (ON142640) possessed variations in 14 places. Nucleotide variations at four (254, 305, 540, 588) and 14 positions (72, 117, 171, 207, 253, 258, 305, 489, 540, 564, 570, 589, 600, 634) when compared with the NCBI accessions U72977 and U72974 for 'C' strain of FAW collected from maize, sweet corn and sorghum crops in India (Swamy *et al.*, 2018). In the study, nucleotide variations were observed in 22 (72, 108, 159, 390, 428, 429, 430, 431, 434, 435, 436, 438, 441, 465, 471, 501, 502, 508, 510, 512, 513 and 516) and 21 (72, 108, 159, 390, 428, 429, 430, 431, 434, 435, 436, 438, 441, 465, 471, 501, 502, 510, 512, 513 and 516) positions in strains, OM462844 (Pearl millet 4 TNAU Coimbatore India) and OM465797 (Pearl millet 5 TNAU Coimbatore India) which are suspected to be 'C' strain populations. Though the strain OM463624 (Pearl millet 2 TNAU Coimbatore India) is grouped in 'R' strain populations, nucleotide variations are found to appear in five places *viz.*, 34, 36, 61, 69 and 84, and hence more genetic distance was ensured. Also, the strain ON142640 (Pearl millet Vridhachalam India) is found to have nucleotide variations in 12 places *viz.*, 72, 108, 159, 381, 390, 445, 459, 465, 468, 486, 487, 512, 513 and 516. The variations in nucleotides may not be due to the host shift but owing to the inter-mating nature of the populations.

### **Phylogenetic tree analysis**

The data matrix included 39 sequences from MtCOI genome of two taxa: one taxon includes 38 FAW sequences, of them 13 across the globe and 25 from India including the test sequences (Fig. 3).

Another taxon, *S. exigua* served as an out group. The phylogenetic tree built using maximum likelihood (ML) following Tamura 3-parameter model showed that *S. frugiperda* divides into two clear clades in which the strain, ON142640 (Pearl millet Vridhachalam India) totally diverged from all the FAW sequences subjected for analysis. The second clade contained most FAW sequences that were clustered and closely related. The rest of the strains are grouped under this clade and are further divided into two clusters. One cluster contained 'R' strain individuals and another comprised 'C' strain individuals. The strains viz. OM462675 (Pearl millet 3 TNAU Coimbatore India), ON114178 (Pearl millet Ramanathapuram India), OM463624 (Pearl millet 2 TNAU Coimbatore India) and ON141589 (Pearl millet Thalavaipuram India) possessed 'R' strain identity whereas, OM462844 (Pearl millet 4 TNAU Coimbatore India) and OM465797 (Pearl millet 5 TNAU Coimbatore India) showed 'C' strain identity. The per cent identity of FAW ranged from 99 to 100. The sequence difference count matrix for the FAW sequences ranged from 0 to 47. The FAW population feeding on sugarcane from Anakapalle formed a separate clade in dendrogram that showed 99.75% resemblance to the Mexico population with 'C' strain identity (Bhavani *et al.*, 2019). Similarly, Chormule *et al.* (2019) claimed that the FAW occurs in sugarcane crops from Kolapur, Maharashtra, assumed 'C' strain identity and close resemblance with USA FAW populations (U72974, U72975 and U72976). In the present investigation also, the two strains viz. OM462844 (Pearl millet 4 TNAU Coimbatore India) and OM465797 (Pearl millet 5 TNAU Coimbatore India) assumed 'C' strain identity and found close resemblance with MT791636 (Panaji Goa India), MT901173 (Kerala India), OM478579 (Ragi 1 Vridhachalam India), OM478578 (Ragi 1 Salem India), MT073266 (Maize Bangladesh), MK318311 (Maize Mexico), KY472248 (Maize Ghana Northern region), KY472251 (Maize Ghana Volta region). The strain, ON142640 (Pearl millet Vridhachalam India) totally diverged from all populations subjected to phylogenetic analysis in this study. This particular sequence has formed a separate clade and could be suspected as an inter-strain hybrid (R/C).

## **Conclusion**

The 'R' strain has colonized maize, sorghum, sweet corn, and many other crops in India, while the 'C' strain has begun to adapt sugarcane, finger millet and pearl millet. As a result, both groups migrated to India and colonized a variety of crops. Because both strains are interbreeding populations, hybridized strains (R/C or C/R) could emerge, feeding on a wide range of hosts and causing damage to a variety of agricultural crops.

## **Funding and acknowledgements**

Financial assistance by the Council of Scientific and Industrial Research, Ministry of Science and Technology, Government of India (File No. 09/641(0175)/2020-EMR-I) to Mr. T. Sathyan (CSIR Direct SRF) is greatly acknowledged. We sincerely acknowledge the financial support given by the Government of Tamil Nadu (GoTN) (Scheme No. F36OT).

### **Ethics statement**

No specific permits were required for the described field studies because no human or animal subjects were involved in this research

### **Originality and plagiarism**

All the data in this manuscript are our original works. This work is not copied from any other article and has not been published anywhere.

### **Consent for publication**

All the authors agreed to publish the content

### **Competing interests**

There was no conflict of interest in the publication of this content

### **Data availability**

All the data of this manuscript are included in the MS.

### **Author contributions**


Idea conceptualization-NS, Experiments- TS, TS, NS, EK, Guidance -NS, SM, VB, EK, JSK, RR, Writing original draft - TS, NS, SM, VB, EK, JSK, RR, TS, Writing- reviewing & editing - TS, NS, SM, VB

### **References**


- Assefa, Y. 2019. Molecular identification of the invasive strain of *Spodoptera frugiperda* (JE smith) (Lepidoptera: Noctuidae) in Swaziland. *Int. J. Trop. Insect Sci.*, **39**: 73–78. <https://doi.org/10.1007/s42690-019-00018-5>.
- Bhavani, B., Sekhar, C.V., Varma, K.P., Lakshmi, B.M., Jamuna, P. and Swapna, B. 2019. Morphological and molecular identification of an invasive insect pest, fall armyworm, *Spodoptera frugiperda* occurring on sugarcane in Andhra Pradesh, India. *J. Entomol. Zool. Stud.*, **7**(4): 12-18
- Casmuz, A., Juarez, M.L., Socias, M.G., Murua, M.G., Prieto, S., Medina, S., Willink, E. and Gastaminza, G. 2010. *Revision de los hospederos del gusano cogollero del maíz, Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Revista de la Sociedad Entomológica Argentina*, **69**: 209–231.

- Chormule, A., Shejawal, N., Nagol, J. and Brown, M.E. 2019. First report of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera, Noctuidae) on sugarcane and other crops from Maharashtra, India. *J. Entomol. Zool. Stud.*, **7**: 114–117.
- Directorate of Millets Development. 2020. Available online at: <http://millets.dacfw.nic.in>.
- Doyle, J.J. and Doyle, J.L. 1990. Isolation of plant DNA from fresh tissue. *Focus*. **12**: 13–15.
- ICAR-IIMR Annual Report. 2019. ICAR-Indian Institute of Maize Research Punjab Agricultural University Campus, Ludiana - 141 004.
- Mauro, D.S., Gower, D.J., Zardoya, R. and Wilkinson, M. A. 2006. Hotspot of gene order rearrangement by tandem duplication and random loss in the vertebrate mitochondrial genome. *Mol. Biol. Evol.*, **23**: 227.
- Montezano, D.G., Specht, A., Sosa-Gómez, D.R., Roque-Specht, V.F., Sousa-Silva, J.C., Paula Moraes, S.V., Peterson, J.A. and Hunt, T.E. 2018. Host plants of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Americas, *Afr. Entomol.*, **26**(2): 286–300.
- Nagoshi, R.N., Goergen, G. and Tounou, K.A. 2018. Analysis of strain distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-Saharan Africa. *Sci. Rep.*, **8**: 3710. <https://doi.org/10.1038/s41598-018-21954-1>.
- Nagoshi, R. N. and Meagher, R. L. 2016. Using intron sequence comparisons in the triose-phosphate isomerase gene to study the divergence of the fall armyworm host strains. *Insect Mol. Biol.*, **25**: 324–337. <https://doi.org/10.1111/imb.12223>.
- Nagoshi, R. N., Silvie, P., Meagher, R. L., Lopez, J. and Machado, V. 2007. Identification and comparison of fall armyworm (Lepidoptera: Noctuidae) host strains in Brazil, Texas, and Florida. *Ann. Entomol. Soc. Am.*, **100**(3): 394-402.
- Nagoshi, R.N. and Meagher, R. 2003. FR tandem-repeat sequence in (Lepidoptera: Noctuidae) host strains. *Ann. Entomol. Soc. Am.*, **96**: 329–335.
- Nagoshi, R.N. and Meagher, R.L. 2004. Behavior and distribution of the two fall armyworm host strains in Florida. *Fla. Entomol.*, **87**(4): 440–9.
- Pashley, D. P. 1998. Quantitative genetics, development, and physiological adaptation in host strains of fall armyworm. *Evol.*, **42**: 93-102.
- Prowell, D. P., McMichael, M. and Silvain, J. F. 2004. Multilocus genetic analysis of host use, introgression, and speciation in host strains of fall armyworm (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **97**(5): 1034-1044.

- Reddy, S.P., Satyavathi, C.T., Khandelwal, V., Patil, H.T., Gupta, P.C., Sharma, L.D., Mungra, K.D., Singh, S.P., Narasimhulu, R., Bhadarge, H.H., Iyanar, K., Tripathi, M.K., Yadav, D., Bhardwaj, R., Talwar, A.M., Tiwari, V.K., Kachole, U.G., Sravanti, K., Priya, S. M., Athoni, B.K., Anuradha, N., Govindaraj, M., Nepolean, T. and Tonapi, V.A. 2021. Performance and Stability of Pearl Millet Varieties for Grain Yield and Micronutrients in Arid and Semi-Arid Regions of India. *Front. Plant Sci.*, **12**: 670201. doi: 10.3389/fpls.2021.670201.
- Sartiami, D., Dadang, Harahap, I.S., Kusumah, Y.M. and Anwar, R. 2020. First record of fall armyworm (*Spodoptera frugiperda*) in Indonesia and its occurrence in three provinces. IOP Conf. Series: *Environ. Earth Sci.*, **468**: 012-021. doi:10.1088/1755-1315/468/1/012021.
- Season and Crop Report 2022. Season and Crop Report of Tamil Nadu 2020-21. FASLI-1430. Department of Economics and Statistics, Chennai – 6. <https://www.tn.gov.in/crop/>
- Sharma, H.C., Davies, J.C. and Arora, J. 1981. Bibliography of insect and non-insect animal pests of millets during 1914-1980. ICRISAT Sorghum and Millets Information Centre, Patancheru, A.P., India.
- Swamy, H. M. M., Asokan, R, Kalleshwaraswamy, C.M., Sharanabasappa, Prasad, Y.G.M., Maruthi, M.S. and Shashank, P.R. 2018. Prevalence of “R” strain and molecular diversity of fall army worm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in India. *Indian J. Entomol.*, **80**: 544.
- Tamura, K., Stecher, G. and Kumar, S. 2021. MEGA 11: Molecular Evolutionary Genetics Analysis Version 11. *Mol. Biol. Evol.*, <https://doi.org/10.1093/molbev/msab120>.
- Verma, S.K. 1980. Field pests of pearl millet. *Trop. Pest Manag.*, **26**: 13-20.



**Fig. 1A** Agarose gel displaying PCR products of FAW collected from pearl millet



**Fig. 1B** Strain-specific RFLPs from the mitochondrial *COI* gene *SacI* digestion



**Fig. 1C** Strain-specific RFLPs from the mitochondrial *COI* gene *MspI* digestion

Lane L– Ladder (100 bp); Lanes 1-5 – TNAU, Coimbatore; Lanes 6-10 – KVK, Ramanathapuram;  
Lanes 11-15 – Thalavaipuram, Virudhunagar; Lanes 16-20 – RRS, Vridhachalam

**Table 1. Details on the FAW collected from pearl millet in different locations of Tamil Nadu**

<b>S. No.</b>	<b>Location</b>	<b>District</b>	<b>Life stage</b>	<b>Sample size</b>	<b>Date of collection</b>	<b>GPS information</b>	<b>NCBI Accession number(s) received</b>
1.	New Area, TNAU	Coimbatore	Four Larvae and one pupa	05	05.10.2019	Latitude: 11.07 N Longitude: 76.99 E	OM462844, OM465797, OM462675, OM463624
2.	KVK, Ramanathapuram	Ramanathapuram	Larvae	05	24.09.2020	Latitude: 11.53 N Longitude: 79.34 E	ON114178
3.	Thalavaipuram	Virudhunagar	Larvae	05	24.09.2020	Latitude: 9.39 N Longitude: 77.50 E	ON141589
4.	RRS, Vridhachalam	Cuddalore	Larvae	05	24.09.2020	Latitude: 11.53 N Longitude: 79.34 E	ON142640

		34	36		61	69	72		84	
OM462844-Pearlmillet4-TNAU-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
OM465797-Pearlmillet5-TNAU-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
ON142640-Pearlmillet1-Vridhachalam-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
OM463624-Pearlmillet2-TNAU-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
ON141589-Pearlmillet1-Thalavaipuram-India	--TTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	58	
ON114178-Pearlmillet1-Ramanathapuram-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
OM462675-Pearlmillet3-TNAU-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
MT605970-Sugarcane-Coimbatore-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
MH753326-Maize-Shivamogga-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
MH704433-Maize-Chikballapur-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	
MH88153-Hassan-India	CTTTA	TT	GGAGATGATCAAA	TTTATAA	TACTTTG	TAACTG	TCATGCTTTT	ATAA	88	

		108						
OM462844-Pearlmillet4-TNAU-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
OM465797-Pearlmillet5-TNAU-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
ON142640-Pearlmillet1-Vridhachalam-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
OM463624-Pearlmillet2-TNAU-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
ON141589-Pearlmillet1-Thalavaipuram-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
ON114178-Pearlmillet1-Ramanathapuram-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
OM462675-Pearlmillet3-TNAU-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
MT605970-Sugarcane-Coimbatore-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
MH753326-Maize-Shivamogga-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
MH704433-Maize-Chikballapur-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148
MH88153-Hassan-India	TTTTTTT	TATAGTTATACC	ATTATAA	TGGAGG	ATTGGAA	TGACTTG	TACCTTTAA	148

		159					
OM462844-Pearlmillet4-TNAU-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
OM465797-Pearlmillet5-TNAU-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
ON142640-Pearlmillet1-Vridhachalam-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
OM463624-Pearlmillet2-TNAU-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
ON141589-Pearlmillet1-Thalavaipuram-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
ON114178-Pearlmillet1-Ramanathapuram-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
OM462675-Pearlmillet3-TNAU-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
MT605970-Sugarcane-Coimbatore-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
MH753326-Maize-Shivamogga-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
MH704433-Maize-Chikballapur-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208
MH88153-Hassan-India	TATTAGGAGC	CCTGATATAGCT	TTCCCACGTATAA	ATAAATAT	AAGTTTTG	GACTTTTAC	208


		381						
OM462844-Pearlmillet4-TNAU-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
OM465797-Pearlmillet5-TNAU-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
ON142640-Pearlmillet1-Vridhachalam-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
OM463624-Pearlmillet2-TNAU-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
ON141589-Pearlmillet1-Thalavaipuram-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
ON114178-Pearlmillet1-Ramanathapuram-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
OM462675-Pearlmillet3-TNAU-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
MT605970-Sugarcane-Coimbatore-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
MH753326-Maize-Shivamogga-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
MH704433-Maize-Chikballapur-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388
MH88153-Hassan-India	CTATTTC	CACCTTCAT	TTAGCTGGAA	TTTCAT	CTATT	TAGGAGCT	ATTAA	388

	390	428-431	434-436	438	441	445		
OM462844-Pearlmillet4-TNAU-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
OM465797-Pearlmillet5-TNAU-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
ON142640-Pearlmillet1-Vridhachalam-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
OM463624-Pearlmillet2-TNAU-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
ON141589-Pearlmillet1-Thalavaipuram-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
ON114178-Pearlmillet1-Ramanathapuram-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
OM462675-Pearlmillet3-TNAU-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
MI605970-Sugarcane-Coimbatore-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
MH753326-Maize-Shivamogga-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
MH704433-Maize-Chikballapur-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
MH88153-Hassan-India	C	ACTATTATTAATATACGATTA	AAATAATTATCATTG	CAAA	CA	TTT	TTT	448
	*	*****	*****	*	*	*	*	*

	459	465	468	471	486,487	501,502	508
OM462844-Pearlmillet4-TNAU-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
OM465797-Pearlmillet5-TNAU-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
ON142640-Pearlmillet1-Vridhachalam-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
OM463624-Pearlmillet2-TNAU-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
ON141589-Pearlmillet1-Thalavaipuram-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
ON114178-Pearlmillet1-Ramanathapuram-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
OM462675-Pearlmillet3-TNAU-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
MI605970-Sugarcane-Coimbatore-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
MH753326-Maize-Shivamogga-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
MH704433-Maize-Chikballapur-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
MH88153-Hassan-India	GAGCTGIAGC	AATTAC	SCATTT	TTATTATTATTATC	TACCTGTTTTAGC	GAGCTT	508
	*	*	*	*	*	*	*

	510	512,513	516		
OM462844-Pearlmillet4-TNAU-India	TAA	ATTT	TTACTTACTGATCGAAATTTAAATACATC	TTTTTGGATCC-----	587
OM465797-Pearlmillet5-TNAU-India	TAA	ATTT	TTACTTACTGATCGAAATTTAAATACATC	TTTTTGGATCCCT-----	590
ON142640-Pearlmillet1-Vridhachalam-India	TAA	ATTT	TTACTTACTGATC-----	-----	529
OM463624-Pearlmillet2-TNAU-India	TAA	ATTT	TTACTTACTGATCGAAATCTAAATACATC	TTTTTGGATCCCT-----	590
ON141589-Pearlmillet1-Thalavaipuram-India	TAA	ATTT	TTACTTACTGATCGAAATCTAAATACATC	TTTTTGGATCCCTGCAGGAGG--	538
ON114178-Pearlmillet1-Ramanathapuram-India	TAA	ATTT	TTACTTACTGATCGAAATCTAAATACATC	TTTTTGGATCCCTGCAGGAGG--	594
OM462675-Pearlmillet3-TNAU-India	TAA	ATTT	TTACTTACTGATCGAAATCTAAATACATC	TTTTTGGATCCCT-----	590
MI605970-Sugarcane-Coimbatore-India	TAA	ATTT	TTACTTACTGATC-----	-----	529
MH753326-Maize-Shivamogga-India	TAA	ATTT	TTACTTACTGATC-----	-----	529
MH704433-Maize-Chikballapur-India	TAA	ATTT	TTACTTACTGATC-----	-----	529
MH88153-Hassan-India	TAA	ATTT	TTACTTACTGATC-----	-----	529
	*	*	*	*	

**Fig.2. Nucleotide variations in the test and other FAW sequences**



**Fig.3 Phylogenetic tree showing the relationships among the FAW strains. Bootstrap support is indicated on the branches**