MANAGEMENT OF THE COFFEE LEAF MINER,

Perileucoptera coffeella (GUÉR-MENV., 1842), IN JAMAICA:

EPIDEMIOLOGY, ECONOMIC IMPORTANCE,

PARASITOID COMPLEX AND EFFECT OF SELECTED

!NSECTICIDES AND THEIR ECOLOGICAL IMPACT

KATHY MARISSA DALIP 1998

## ABSTRACT

Monthly monitoring of the coffee leaf miner (CLM), *Perileucoptera coffeella* (Guérin-Méneville, 1842), was carried out on two coffee farms, one each at Oxford and Mt. Hill, during May 1992 - May 1994, and every four to six months thereafter. The infestation of the CLM fluctuated in a cyclic pattern with peaks in May 1992 (85.6%) and January 1993 (64.9%) at Oxford, and in September (14.1%) and May 1992 (11.4%) in Mt. Hill; the lowest levels were in June 1993 (12.1%) and May 1994 (11.9%) at Oxford, and April 1993 (2.0%) at Mt. Hill. Low infestation followed periods of high rainfall. The population fell between 0 and 5% between July 1994 and March 1998 at the plantations but is increasing again due to El Niño-induced drought.

The mean monthly infestation pattern within plantations at Oxford varied between 39.05 and 45.89%, which was lower in plots bordering the river bank than the central ones and those surrounded by trees of var. *caturra*. In Mt. Hill, infestation in plots was not significantly (P > 0.05) different and ranged between 5.39 and 5.82%.

There was a significant (P < 0.001) difference in the mean infestation of different strata of the trees; at Oxford, it was upper (51.18%) > middle (41.68%) > lower (36.45%) and at Mt. Hill, it was lower (7.97%) > middle (4.27%) > upper (4.16%), respectively. The mean infestation was also significantly (P = 0.008) more in the north (44.48%) = east (43.68%) = south (42.41%) = west (41.84%) at Oxford and north (6.21%) > south

In establishing the economic threshold (ET) for the CLM, the damage index, D - which was the product of the LAD and infestation level- was created. A threshold of 300 for the damage index was used to obtain an ET of 15% LAD at 20% tree infestation.

Fourteen species of parasitoids - two braconids, eight eulophids, two pteromalids, and two other unclassified species - emerged from CLM-infested field-collected leaves. The level of parasitism ranged between 2.44 and 62.86% and varied significantly (P = 0.05) among some plantations.

Decis 2.5EC (deltamethrin), Karate 2.5EC ( $\lambda$  cyhalothrin), Furadan 10G (carbofuran) and Miral 500CS (isazofos) applied by farmers gave no significant (P > 0.05) reduction in the CLM infestation. Field trials with Miral 500CS, Basudin 600 (diazinon), Chlorpyrifos 2E and crude extracts of *Artocarpus altilis* (breadfruit), *Blighia sapida* (ackee) and *Azadirachta indica* (neem) revealed that only Miral and ackee oil had any significant (P < 0.05) toxic effect on larvae of the CLM. None of the insecticides had a significant (P > 0.05) effect on the level of infestation. The feeding activity of the CLM was inhibited by for two weeks by ackee oil and for at least three weeks by neem, Miral and Basudin.

The persistence ( $t_{\frac{1}{2}}$  in days) on coffee leaves and soil (data in parentheses) of test insecticides was chlorpyrifos - 1.1 (5.8); isazofos - 1.3 (3.1); diazinon - 1.8 (7.3); dimethoate - 2.1 (3.2); deltamethrin - 5.9 (1.8);  $\lambda$ 

cyhalothrin - 12.7 (6.5). The insecticides did not have prolonged, adverse effects the soil fauna.

An IPM strategy, integrating EIL with tree pruning, irrigation, fertilization, drought-resistant root stock, parasitoids and plant extracts, is proposed.

deln L. Rabinson whose assistance with emailine, advice,

Processed and especially Mr. K. Onbey - In providing relevant